



## Overstory community composition and elevated atmospheric CO<sub>2</sub> and O<sub>3</sub> modify understory biomass production and nitrogen acquisition

Janet M. Bandeff<sup>1</sup>, Kurt S. Pregitzer<sup>1,3</sup>, Wendy M. Loya<sup>1</sup>, William E. Holmes<sup>2</sup> & Donald R. Zak<sup>2</sup>

<sup>1</sup>*Ecosystem Science Center, School of Forest Resources and Environmental Science, Michigan Technological University, 1400 Townsend Avenue, Houghton, MI, 49931, USA.* <sup>2</sup>*School of Natural Resources and the Environment, and Department Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, 48109-1115, USA.* <sup>3</sup>*Corresponding author\**

Received 27 June 2005. Accepted in revised form 13 December 2005

**Key words:** climate change, global change, light, nitrogen cycling, plant competition, stable isotopes

### Abstract

Elevated atmospheric CO<sub>2</sub> and O<sub>3</sub> have the potential to affect the primary productivity of the forest overstory, but little attention has been given to potential responses of understory vegetation. Our objective was to document the effects of elevated atmospheric CO<sub>2</sub> and O<sub>3</sub> on understory species composition and biomass and to quantify nitrogen (N) acquisition by the understory vegetation. The research took place at the aspen free-air CO<sub>2</sub> and O<sub>3</sub> enrichment (FACE) experiment, which has four treatments (control, elevated CO<sub>2</sub>, elevated O<sub>3</sub>, and elevated CO<sub>2</sub>+O<sub>3</sub>) and three tree communities: aspen, aspen/birch, and aspen/maple. In June 2003, each FACE ring was uniformly labeled with <sup>15</sup>N applied as NH<sub>4</sub>Cl. Understory biomass was harvested in June of 2004 for productivity, N, and <sup>15</sup>N measurements, and photosynthetically active radiation (PAR) was measured below the canopy. The understory was divided into five species groups, which dominate in this young aggrading forest: *Taraxacum officinale* (dandelion), *Solidago* sp. (goldenrod), *Trifolium repens* and *T. pretense* (clover), various species from the *Poaceae* family (grass), and composited minor components (CMC). Understory species composition, total and individual species biomass, N content, and <sup>15</sup>N recovery showed overstory community effects, but the direct effects of treatments was masked by the high variability of these data. Total understory biomass increased with increasing light, and thus was greatest under the open canopy of the aspen/maple community, as well as the more open canopy of the elevated O<sub>3</sub> treatments. Species were different from one another in terms of <sup>15</sup>N recovery, with virtually no <sup>15</sup>N recovered in clover and the greatest amount recovered in dandelion. Thus, understory species composition and biomass appear to be driven by the structure of the overstory community, which is determined by the tree species present and their response to the treatments. However, N acquisition by the understory does not appear to be affected by either the overstory community or the treatments at this point.

### Introduction

Since the start of the Industrial Revolution, atmospheric CO<sub>2</sub> has increased from 280 to 377 μL L<sup>-1</sup> today. Data from free-air carbon

dioxide enrichment (FACE) experiments demonstrate that elevated levels of CO<sub>2</sub> led to greater CO<sub>2</sub> assimilation by plants and increased productivity (Ainsworth and Long, 2005; Norby et al., in press; Nowak et al., 2004). However, increased levels of tropospheric O<sub>3</sub> have also accompanied increases in atmospheric CO<sub>2</sub>, with the amount

\* E-mail: kspregit@mtu.edu

of increase depending on geographic location (Fishman et al., 2003). Elevated  $O_3$  is harmful to plant tissues, net photosynthesis (Reich and Amundson, 1985), and biomass production (Reich 1987). The interaction between  $CO_2$  and  $O_3$  is not well understood, although evidence suggests that elevated  $CO_2$  will alleviate some of the negative effects of elevated  $O_3$ , which include reduced leaf area, leaf and root biomass, net primary productivity, and water-use efficiency (Karnosky et al., 2003).

Elevated  $CO_2$  and  $O_3$  may also alter the cycling of nitrogen (N) within forested ecosystems (McGuire et al., 1995). In order for trees to maintain increased productivity under elevated  $CO_2$ , they must increase N use efficiency or N uptake (Pataki et al., 2003). A study of nutrient cycling at the Oak Ridge National Laboratory FACE experiment (Johnson et al., 2004) provided evidence that trees are able to increase their N uptake, likely through greater fine root production, although the source of this additional N is unclear.

Elevated  $CO_2$  research in forests has focused primarily on woody plants (Poorter and Navas, 2003). Existing research on herbaceous species has mostly been in an agricultural setting, and the forest herbaceous understory has been little studied. While understory vegetation does not contribute a significant amount of biomass to forested ecosystems, it is important for biodiversity and wildlife habitat. Meta-analyses of FACE data suggest that herbaceous plants are less responsive than trees to elevated  $CO_2$ , (Ainsworth and Long, 2005; Nowak et al., 2004), with fast-growing C3 herbs being more responsive than slow-growing C3 herbs or C4 plants (Poorter and Navas, 2003). Herbaceous plants also experience a greater reduction in leaf nitrogen content compared to woody plants (Nowak et al., 2004).

The direct effects of elevated atmospheric  $CO_2$  and  $O_3$  on understory biomass production and N acquisition, as well as the indirect effects of increased overstory biomass and N demand, are not known. Total understory biomass, individual species biomass, and species composition of the understory vegetation are known to be affected by the amount of light reaching the forest floor, which is driven by overstory species composition, as well as stand and crown structure (Alaback, 1982; Berger and Puettmann, 2000; Hunt et al., 2005; Legare et al., 2002; Peek et al., 2001). Understory species

composition and biomass are also affected by leaf litter (Harrington et al., 2003), productivity (Adkison and Gleeson, 2004), soil nutrient concentrations (Hunt et al., 2005), nitrogen deposition rate (Rainey et al., 1999), throughfall precipitation (Anderson et al., 1969), above and below ground competition with overstory vegetation (Harrington et al., 2003; Riegel et al., 1992), disturbance (Moser et al., 1996), and successional dynamics (Gleeson and Tilman, 1990).

The objectives of this research were to study the effects of elevated levels of atmospheric  $CO_2$  and  $O_3$  on understory composition and biomass production and to quantify N acquisition by the understory vegetation using a pulse-chase  $^{15}N$  tracer experiment. We hypothesized that understory species biomass would be lowest in the elevated  $CO_2$  treatment, which is expected to have the lowest light levels, and highest in the elevated  $O_3$  treatment, which is expected to have the highest light levels. We also hypothesized that recovery of the  $^{15}N$  tracer would be lowest in the elevated  $CO_2$  treatments because of greater demand for nitrogen by the overstory trees under elevated  $CO_2$  (Oren et al., 2001).

## Methods

### *Study site*

The Aspen FACE site is located at the Harshaw Experimental Farm outside of Rhinelander, Wisconsin, USA. The 32 hectare field contains 12 FACE rings, which are 30 m in diameter. Three rings are exposed to one of four treatments: control, elevated  $CO_2$ , elevated  $O_3$ , and elevated  $CO_2 + O_3$ . Elevated levels of  $CO_2$  and  $O_3$  are maintained at approximately 150% of ambient levels using a trace gas monitoring system, high-volume blowers, and vertical vent piping (Dickson et al., 2000; Karnosky et al., 2003).  $CO_2$  levels average 356 and 534  $\mu L L^{-1}$  for control and elevated  $CO_2$ , respectively.  $O_3$  levels average 36 and 50 nL  $L^{-1}$  for control and elevated  $O_3$ , respectively.  $O_3$  levels were calculated as 12 h daytime mean  $O_3$  exposure. Detailed hourly  $O_3$  exposure showing maximum daily  $O_3$  concentration can be found in Karnosky et al. (2005). The 12 rings were divided into three blocks, with 1 ring of each treatment composing a block. Each ring was

divided into three communities (aspen, aspen/birch, and aspen/maple) by planting half of the ring with five different trembling aspen clones (*Populus tremuloides* Michx.), a quarter with alternating individuals of trembling aspen and paper birch (*Betula papyrifera* Marsh.), and the final quarter with trembling aspen and sugar maple (*Acer saccharum* Marsh.). The seedlings were planted at 1×1-m spacing in June of 1997. The soil was plowed and tilled before planting. Fumigation of the rings began in spring of 1998 (Dickson et al., 2000). The understory vegetation was controlled in 1997, 1998, and 1999 by applying Roundup two times per year using a backpack sprayer. Following the final herbicide application, the understory vegetation was allowed to develop. An ecosystem-scale <sup>15</sup>N tracer experiment was conducted in June 2003 in which all FACE rings were uniformly labeled with <sup>15</sup>N. The tracer was applied in the form of NH<sub>4</sub>Cl at the rate of 0.0153 g <sup>15</sup>N m<sup>-2</sup> using backpack sprayers. Immediately following <sup>15</sup>N application, the rings were irrigated with 0.2 cm of water to rinse the <sup>15</sup>N solution off the understory plants and into the soil.

The amount of CO<sub>2</sub> and O<sub>3</sub> reaching the forest understory vegetation is not significantly different from that reaching the canopy. Passive samplers installed in an O<sub>3</sub> treatment ring to determine monthly cumulative O<sub>3</sub> show no significant difference between 1 m and canopy samplers (Karnosky, personal communication). Analysis of spatial control of the forest face prototype designed to test the feasibility of using FACE technology to fumigate mature loblolly pine trees found that 3D CO<sub>2</sub> concentrations were acceptably homogenous (Hendrey et al., 1999).

#### *Photosynthetically active radiation*

Because the light reaching the understory layer of the forest is directly related to forest overstory biomass production (Chazdon, 1988), we measured the amount of photosynthetically active radiation (PAR) reaching the understory vegetation. A Sunfleck Ceptometer (Decagon Devices, Inc., Pullman WA) was used to measure PAR at four positions in the aspen community and two positions in the aspen/birch and aspen/maple communities on June 23, 2004, at the same time

the understory was harvested, and again on July 27, 2004, at the time of approximate peak leaf area for the season. Measurements were taken between 11:00 am and 3:00 pm local time on clear, cloudless days. At each position, the ceptometer was held at arms length, level to the ground and below all tree branches, and turned slowly in a circle. Eight measurements were taken along the circle approximately 45° apart and averaged (Burton et al., 1991). One circle was also established and measured in an open area outside of each ring in the manner described above. Two positions were used in the aspen/birch and aspen/maple communities originally because these areas are half the size of the aspen community. For the July measurement period, four positions were sampled in order to obtain better precision.

#### *Understory harvest*

The understory was harvested on June 21–24, 2004, just before the overstory vegetation was sampled for N analysis. The understory was clipped at ground level within four 0.5 m<sup>2</sup> quadrats per community. Quadrat locations were randomly preselected such that, within a community, the quadrats were in the same location in each ring. However, it was necessary to stay away from certain randomly placed equipment and boardwalks throughout the ring. Therefore, the actual quadrat location was the spot nearest to the randomly preselected location that met equipment and boardwalk criteria. Once clipped, the vegetation was sorted into the following fourteen groups: dandelion (*Taraxacum officinale*), goldenrod (*Solidago* sp.), red and white clover (*Trifolium repens* and *T. pretense*), grasses (C3 grasses from the Poaceae family), ferns, tree seedlings, yarrow (*Achillea millefolium*), yellow rocket (*Barbarea vulgaris*), annual fleabane (*Erigeron annuus*), strawberry (*Fragaria* sp.), orange hawkweed (*Hieracium aurantiacum*) and its nearly identical counterpart, prickly lettuce (*Lactuca scariola*), plantain (*Plantago* sp.), and low hop clover (*Trifolium campestre*). Several additional species were seen in the understory, but could not be identified. These were placed in an ‘other’ category. This category never accounted for more than 7% of the biomass in a quadrat and averaged 0.3%. Two samples of dandelion, goldenrod, clover, and grass were also taken in similar habitats

outside of the rings to determine the natural abundance of  $^{15}\text{N}$ . These species were chosen because they represented over 90% of the total understory biomass.

#### *Laboratory and statistical analyses*

The understory samples were dried in a 65 °C oven for 72 h and weighed to the nearest 0.1 g. The samples were then composited by community into five groups: dandelion, goldenrod, clover, grass, and composited minor components (CMC). All composite samples were ground into a powder using a ball mill. Approximately 3 mg of ground sample were weighed into a tin capsule and analyzed for N concentration and  $\delta^{15}\text{N}$  using a Costech Elemental Combustion System connected to a Thermo Finnigan ConFloIII Interface and Delta<sup>plus</sup> Continuous Flow-Stable Isotope Mass Spectrometer. International Atomic Energy Agency, United States Geological Survey, and National Institute of Standards and Technology certified isotopic standards were run at the beginning of every analysis. Every ten samples were followed by an internal standard and every 25th sample was an analytical replicate. N concentration and  $\delta^{15}\text{N}$  were used to calculate N content,  $^{15}\text{N}$  atom percent excess, and  $^{15}\text{N}$  recovery.

Statistical analysis of understory biomass, N concentration, N content, atom percent excess  $^{15}\text{N}$ , and  $^{15}\text{N}$  recovery was performed in SAS (version 8.2, SAS Institute Inc., Cary, NC) using PROC GLM for a split-plot randomized complete block design as detailed in King et al. (2001). An alpha level of 0.05 was used and significant differences were further analyzed using least squared means. Due to uneven variance in the biomass estimates under elevated  $\text{CO}_2$  and  $\text{O}_3$ , we multiplied the biomass values by 10 and log transformed these data. Statistical results were similar to the original analysis. Therefore, the results presented below are for untransformed data.

## **Results**

### *Photosynthetically active radiation*

The amount of PAR reaching the understory was significantly different among the three communities, with the amount of PAR reaching the

understory being 30 and 60% lower in the aspen/birch community than the aspen and aspen/maple communities, respectively ( $P < 0.001$ ). In the aspen and aspen/birch communities,  $\text{CO}_2$  significantly decreased PAR (main effect  $P = 0.04$ ) and  $\text{O}_3$  significantly increased PAR (main effect  $P = 0.02$ ; Figure 1). There were no significant treatment effects in the aspen/maple community.

### *Species composition and biomass*

The understory was dominated by perennial old-field vegetation. More than 30 herbaceous species, 2 fern species, and 5 tree species were found in the understory layer. Dandelion, goldenrod, clover, and grass dominated, accounting for over 90% of the total biomass.

Dandelion, clover, and grass biomass differed significantly among overstory communities ( $P = 0.02$ ,  $< 0.0001$ , and  $0.03$  respectively; Table 1). We found no  $\text{CO}_2$  or  $\text{O}_3$  main effects or  $\text{CO}_2 \times \text{O}_3$  interaction effects on individual species biomass and total understory biomass. However, some treatment trends were apparent. Simple linear regression analysis showed that mean understory biomass and mean PAR by treatment for the aspen and aspen/birch communities showed a positive correlation (data not shown,  $R^2 = 0.69$ ,  $P = 0.01$ ). Additionally, clover biomass increased in treatments exposed to  $\text{O}_3$  compared to those at ambient or elevated  $\text{CO}_2$ , with the increase ranging from 80–430% in the aspen community, 180–400% in the aspen/birch community, and 30–100% in the aspen/maple community. In the aspen and aspen/birch communities, total understory biomass was highest in the  $\text{O}_3$  treatment, while the  $\text{CO}_2 + \text{O}_3$  treatment showed a marked increase in biomass over the control treatment (data not shown).

### *Nitrogen acquisition*

Natural abundance  $\delta^{15}\text{N}$  values were  $-1.5\text{‰}$  for clover,  $-0.71\text{‰}$  for dandelion,  $-0.29\text{‰}$  for goldenrod,  $0.33\text{‰}$  for grass, and  $-0.22\text{‰}$  for CMC. The natural abundance  $\delta^{15}\text{N}$  value of CMC was calculated by taking the average  $\delta^{15}\text{N}$  of dandelion, goldenrod, and grass. N concentration differed by species group with clover having the highest N concentration, but N concentration did




Figure 1. Photosynthetically active radiation under the overstory canopy at the Aspen FACE site on July 27, 2004. Error bars represent one standard error of the mean ( $n=3$ ). Different letters within a community represent significant differences at an alpha level of 0.05.

not differ among treatments or communities. N content differed in dandelion, clover, and grass between overstory communities (Table 1). 15N recovery varied significantly by species group and within species group by overstory community. Clover recovered 96% less 15N than dandelion in the aspen and aspen/birch communities and 83% less in the aspen/maple community. Clover also recovered over 85% less 15N than goldenrod across all communities (Table 1). No significant treatment differences were found.

## Discussion

### *Species composition and biomass*

Understory biomass was on average 3% of total aboveground biomass (aboveground biomass estimates can be found in King et al., in press). Understory biomass appears to be driven by the amount of light reaching the understory vegetation. We had anticipated that the differences in PAR by treatment would have been strong enough to produce significant treatment effects on total and individual species biomass, but this was not the case. Rather, individual species in our

study responded to the overstory community, with clover and grass thriving in the aspen/maple community and dandelion thriving in the aspen/birch community. This suggests that clover may out compete dandelion under open canopies or that dandelion is more shade tolerant than clover. There is no indication in our results that the response of the understory to the overstory community is due to differences in N competition among the different overstory communities, as there were no overstory community effects on understory N concentration or atom percent excess 15N.

The individual species responses to light differences among the overstory communities are similar to the nonsignificant species responses to light seen among the treatments. Understory species composition and biomass varied widely between quadrats and FACE rings, resulting in high standard errors in total and individual species biomass. This heterogeneity many have contributed to the lack of treatment response. In the future, a greater sampling intensity may be necessary in order to determine treatment responses.

Besides sampling intensity, several other factors may have contributed to a lack of biomass response to the elevated CO<sub>2</sub> and O<sub>3</sub> treatments. While PAR is significantly different among the

Table 1. Mean biomass, N content, and <sup>15</sup>N recovery for understory species groups at the Aspen FACE site

	Biomass (g m <sup>-2</sup> )			N Content (gN m <sup>-2</sup> ) <sup>a</sup>			<sup>15</sup> N Recovery (μg <sup>15</sup> N m <sup>-2</sup> ) <sup>b</sup>		
	Aspen	Aspen/Birch	Aspen/Maple	Aspen	Aspen/Birch	Aspen/Maple	Aspen	Aspen/Birch	Aspen/Maple
Clover	15.6 ab (4.0)	9.7 b (4.7)	19.6 a (3.9)	0.44 ab (0.11)	0.29 b (0.13)	0.54 a (0.12)	11.3 (2.3)	13.9 (4.3)	21.1 (6.2)
Dandelion	20.1 b (2.3)	29.5 a (2.6)	12.3 c (1.6)	0.32 b (0.04)	0.53 a (0.06)	0.19 b (0.02)	184.7 b (26.9)	371.4 a (67.4)	127.8 b (17.7)
Goldenrod	15.6 (4.0)	11.1 (3.0)	23.4 (6.4)	0.31 (0.09)	0.24 (0.06)	0.45 (0.1)	92.2 (19.0)	93.6 (28.5)	192 (56.8)
Grass	1.2 b (0.6)	1.4 b (0.6)	4.2 a (1.4)	0.020 b (0.01)	0.026 b (0.01)	0.060 a (0.02)	10.2 b (5.2)	13.0 b (4.6)	33.9 a (9.9)
CMC	4.9 (1.2)	3.1 (0.7)	4.1 (1.4)	0.086 (0.02)	0.066 (0.01)	0.075 (0.03)	47.5 (10.7)	42.9 (8.0)	51.4 (17.0)
Total	57.4 (5.9)	54.8 (8.5)	63.6 (10.1)	1.15 (0.2)	1.11 (0.2)	1.3 (0.2)	328.9 (37.4)	525.8 (74.4)	415.6 (67.3)

Values in parenthesis are one standard error of the mean ( $n = 12$ ). Different letters within a species group represent significant differences at an alpha level of 0.05.

<sup>a</sup>Nitrogen content: biomass multiplied by N concentration.

<sup>b</sup><sup>15</sup>N recovery: N content multiplied by atom percent excess <sup>15</sup>N.

four treatments, the vast majority of the plants found growing in the understory were shade intolerant species typically found in open fields, not forested understories. Also, the large number of factors that affect understory biomass production makes it difficult to tease apart responses to specific treatments. Differing species responses to environmental and community factors affected the total understory biomass response to the treatments. Clover seemed to flourish with elevated O<sub>3</sub>. This could be attributed to several causes, including a direct positive response to O<sub>3</sub>, a response to negative growth by other species because of O<sub>3</sub>, or a response to greater light availability. Environmental and individual species responses were also considered to be the factors driving understory biomass at the Oak Ridge National Laboratory FACE experiment (Belote et al., 2004).

Additionally, under low light conditions, the relative rate of photosynthesis is expected to increase under elevated CO<sub>2</sub> relative to ambient CO<sub>2</sub> (Long and Drake, 1991). Osborne et al. (1997) documented greater carbon uptake in understory Indiana strawberry when grown under elevated CO<sub>2</sub> (*Duchesnea indica*). Hattenschwiler and Korner (1997) found that when grown under elevated CO<sub>2</sub>, shade-intolerant species were able to obtain greater biomass and grow under low light. In our study, the total biomass was higher, although not significantly different, in the CO<sub>2</sub> + O<sub>3</sub> treatment compared to the control, despite similar light levels, and biomass was the same in the control and CO<sub>2</sub> treatments, despite significantly lower light levels in the CO<sub>2</sub> treatment, possibly indicating that understory plants were able to obtain more photosynthate for growth under elevated CO<sub>2</sub>.

The different responses of the species groups to the treatments and communities are to be expected, as our species groups were different in life form and life history attributes. Together these differences between species and their responses to overstory species composition, light, interspecific competition, CO<sub>2</sub>, and O<sub>3</sub> make it difficult to predict community- and ecosystem-level responses to a changing atmosphere. It is, however, apparent that elevated CO<sub>2</sub> and O<sub>3</sub> influence light transmission through the canopy (Figure 1), and this initiates a series of complex interactions in the forest understory.

### *Nitrogen acquisition*

On average, 3% of the  $^{15}\text{N}$  tracer was recovered by the understory vegetation, compared to only 8% in the forest overstory. The majority of the tracer was recovered by the leaf litter and soil organic matter (Holmes, unpublished data). These results suggest that new N from the tracer was available for uptake throughout the growing season as leaf litter turned over. The total amount of N acquired by the understory vegetation, as measured by N content and  $^{15}\text{N}$  recovery, was not significantly affected by the treatments. Other elevated  $\text{CO}_2$  experiments have also found that overstory and understory vegetation are able to obtain approximately equal amounts of N in control and elevated  $\text{CO}_2$  treatments (Finzi et al., 2002; Johnson et al., 2004). Our results demonstrate that, at least in the understory, elevated levels of  $\text{CO}_2$  and  $\text{O}_3$  have not influenced total N acquisition.

N content and  $^{15}\text{N}$  recovery of dandelion and grass differed by overstory community. These differences are likely driven by biomass, as there were no community effects on N concentration and  $^{15}\text{N}$  atom percent excess. However, due to differences in understory species composition between the three overstory communities, total understory biomass was highest in the aspen/maple community, while total  $^{15}\text{N}$  recovery was highest in the aspen/birch community. This is largely due to the fact that dandelion dominated the aspen/birch community, while clover, which recovered very little  $^{15}\text{N}$ , along with goldenrod, dominated the aspen/maple community. Clover likely obtained the majority of its N through biological fixation, rather than through uptake of inorganic N, and therefore acquired little of the  $^{15}\text{N}$  tracer.

Individual species N content and  $^{15}\text{N}$  recovery did not differ significantly by treatment. Other studies addressing the N content of individual species have found a variety of responses to elevated  $\text{CO}_2$ . A review of FACE data by Nowak et al. (2004) found that all but one herbaceous species studied had reduced leaf N content under elevated  $\text{CO}_2$ . However, a study of understory vegetation in a sweetgum plantation found four of five understory taxa showed no significant change in C:N ratios when grown at ele-

vated  $\text{CO}_2$  (Sanders et al., 2004). Even though N content was unaffected by elevated  $\text{CO}_2$  and  $\text{O}_3$ , species differences in N concentration and changes in species composition with overstory community and treatment could affect N distribution within the understory and perhaps the amount of N remaining in the soil.

In conclusion, overstory community composition, light, and interspecific competition affected species composition and biomass. The effects of the elevated  $\text{CO}_2$  and  $\text{O}_3$  treatments were manifested indirectly through overstory community response. Further, total N content and total  $^{15}\text{N}$  recovery in the understory were unaffected by  $\text{CO}_2$  and  $\text{O}_3$  or community type. However, N concentration and  $^{15}\text{N}$  recovery did differ by understory species groups with clover recovering virtually none of the  $^{15}\text{N}$  tracer regardless of treatment. Elevated  $\text{CO}_2$  and  $\text{O}_3$  appear to have little effect on the understory component of the N cycle in this experiment. Nevertheless, changes in the atmosphere did appear to contribute to alterations in species composition of the understory and responses were specific to groups of herbaceous taxa, demonstrating both direct and indirect responses of the forest understory to changes in the Earth's atmosphere.

### **Acknowledgements**

This research was supported by the US Department of Energy's Office of Science (PER: Program for Ecosystem Research), the USDA Forest Service (Northern Global Change Program and North Central Research Station), the National Science Foundation (DEB, DBI/MRI). G. Hendrey, K. Lewin, J. Nagey, D. Karnosky, and J. Sober have been instrumental in the successful implementation of this long-term study.

### **References**

- Adkison G P and Gleeson S K 2004 Forest understory vegetation along a productivity gradient. *J. Torrey Bot. Soc.* 131, 32–44.
- Ainsworth E A and Long S P 2005 What have we learned from 15 years of free-air  $\text{CO}_2$  enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising  $\text{CO}_2$ . *New Phytol.* 165, 351–372.

- Alaback P B 1982 Dynamics of understory biomass in sitka spruce-western hemlock forests of southeast Alaska. *Ecology* 63, 1932–1948.
- Anderson R C, Loucks O L and Swain A M 1969 Herbaceous response to canopy cover, light intensity, and throughfall precipitation in coniferous forests. *Ecology* 50, 255–263.
- Belote R T, Welzin J F and Norby R J 2004 Response of an understory plant community to elevated [CO<sub>2</sub>] depends on differential responses of dominant invasive species and is mediated by soil water availability. *New Phytol.* 161, 827–835.
- Berger A L and Puettmann K J 2000 Overstory composition and stand structure influence herbaceous plant diversity in the mixed aspen forest of northern Minnesota. *Am. Midl. Nat.* 143, 111–125.
- Burton A J, Pregitzer K S and Reed D D 1991 Leaf area and foliar biomass relationships in northern hardwood forests located along an 800-km acid deposition gradient. *Forest Sci.* 37, 1041–1059.
- Chazdon R L 1988 Sunflecks and their importance to forest understorey plants. *Adv. Ecol. Res.* 18, 1–63.
- Dickson R E, Lewin K F, Isebrands J G, Coleman M D, Heilman W E, Riemenschneider D E, Sober J, Host G E, Zak D R, Hendrey G R, Pregitzer K S and Karnosky D F 2000 Forest atmosphere carbon transfer and storage (FACTS-II) – the aspen free-air CO<sub>2</sub> and O<sub>3</sub> enrichment (FACE) project: an overviews Technical Report NC-214. Washington DC, USDA.
- Finzi A C, DeLucia E H, Hamilton J G, Richter D D and Schlesinger W H 2002 The nitrogen budget of a pine forest under free air CO<sub>2</sub> enrichment. *Oecologia* 132, 567–578.
- Fishman J, Wozniak A E and Creilson J K 2003 Global distribution of tropospheric ozone from satellite measurements using the empirically corrected tropospheric ozone residual technique: identification of the regional aspects of air pollution. *Atmos. Chem. Phys. Discuss.* 3, 1453–1476.
- Gleeson S K and Tilman D 1990 Allocation and the transient dynamics of succession on poor soils. *Ecology* 71, 1144–1155.
- Harrington T B, Dagley C M and Edwards M B 2003 Above- and below-ground competition from longleaf pine limits performance of reintroduced herbaceous species. *Forest Sci.* 49, 681–695.
- Hattenschwiler S and Korner C 1997 Growth of autotrophic and root-hemiparasitic understory plants under elevated CO<sub>2</sub> and increased N deposition. *ACTA Oecol.* 18, 327–333.
- Hendrey G R, Ellsworth DS, Lewin K F and Nagy J 1999 A free-air enrichment system for exposing tall forest vegetation to elevated atmospheric CO<sub>2</sub>. *Globel. Change Biol.* 5, 293–309.
- Hunt S L, Gordon A M and Morris D M 2005 Aspects of ecological development in managed stands of jack pine and black spruce in northern Ontario: understory vegetation and nutrient relations. *Forest. Chron.* 81, 61–72.
- Johnson D W, Cheng W, Joslin J C, Norby R J, Edwards N T and Todd D E 2004 Effects of elevated CO<sub>2</sub> on nutrient cycling in a sweetgum plantation. *Biogeochemistry* 69, 379–403.
- Karnosky D F, Pregitzer K S, Zak D R, Kubiske M E, Hendrey G R, Weinstein D, Nosal M and Percy K E 2005 Scaling ozone responses of forest trees to the ecosystem level in a changing climate. *Plant Cell Environ* 28, 965–981.
- Karnosky D F, Zak D R, Pregitzer K S, Awmack C S, Bockheim J G, Dickson R E, Hendrey G R, Host G E, King J S, Kopper B J, Kruger E L, Kubiske M E, Lindroth R L, Mattson W J, McDonald E P, Noormets A, Oksanen E, Parsons W F J, Percy E K, Podila G L, Riemenschneider D E, Sharma P, Thakur R C, Sober A, Sober J, Jones W S, Anttonen S, Vapaavuori E, Mankovska B, Heilman W E and Isebrands J G 2003 Tropospheric O<sub>3</sub> moderates responses of temperate hardwood forests to elevated CO<sub>2</sub>: a synthesis of molecular to ecosystem results from the Aspen FACE project. *Funct. Ecol.* 17, 289–304.
- King J S, Pregitzer K S, Kubiske M E, Hendrey G R, Giardina C P, McDonald E P, Quinn V S and Karnosky D F 2001 Tropospheric O<sub>3</sub> compromises net primary production in young stands of trembling aspen, paper birch and sugar maple in response to elevated atmospheric CO<sub>2</sub>. *New Phytol.*, 168, 623–636.
- King J S, Pregitzer K S, Zak D R, Sober J, Isebrands J G, Dickson R E, Hendrey G R and Karnosky D F 2001 Fine-root biomass and fluxes of soil carbon in young stands of paper birch and trembling aspen as affected by elevated atmospheric CO<sub>2</sub> and tropospheric O<sub>3</sub>. *Oecologia* 128, 237–250.
- Legare S, Bergeron Y and Pare D 2002 Influence of forest composition on understory cover in boreal mixedwood forests in western Quebec. *Silva Fenn.* 36, 353–366.
- Long S P and Drake B G 1991 Effect of the long-term elevation of CO<sub>2</sub> concentration in the field on the quantum yield of photosynthesis of the C3 sedge *Scirpus olneyi*. *Plant Physiol.* 96, 221–226.
- McGuire D A, Melillo J M and Joyce L A 1995 The role of nitrogen in response of forest net primary production to elevated atmospheric carbon dioxide. *Annu. Rev. Ecol. Syst.* 26, 473–503.
- Moser W K, Ducey M J and Ashton P M S 1996 Effects of fire intensity on competitive dynamics between red and black oaks and mountain laurel. *North. J. Appl. For.* 13, 119–123.
- Norby R J, DeLucia E H, Gielen B, Calfapietra C, Giardina C P, King J S, Ledford J, McCarthy H R, Moore D J P, Ceulemans R, De Angelis P, Finzi A C, Karnosky D F, Kubiske M E, Lukac M, Pregitzer K S, Scarascia-Mugnozza G E, Schlesinger W H and Oren R 2005 Forest response to elevated CO<sub>2</sub> is conserved across a board range of productivity. *P. Natl. Acad. Sci-Biol.* 102, 18052–18056.
- Nowak R S, Ellsworth S D and Smith S D 2004 Functional responses of plants to elevated atmospheric CO<sub>2</sub>—do photosynthetic and productivity data from FACE experiments support early predictions?. *New Phytol.* 162, 253–280.
- Oren R, Ellsworth D S, Johnsen K S, Phillips N, Ewers B E, Maier C, Schafer K V R, McMarthy H, Hendrey G, NeNulty S G and Katul G G 2001 Soil fertility limits carbon sequestration by forest ecosystems in a CO<sub>2</sub>-enriched atmosphere. *Nature* 411, 469–472.
- Osborne C P, Drake B G, LaRoche J and Long S P 1997 Does long-term elevation of CO<sub>2</sub> concentration increase photosynthesis in forest floor vegetation? Indiana strawberry in a Maryland forest. *Plant Physiol.* 114, 337–344.
- Pataki D E, Ellsworth D S, Evans R D, Gonzalez-Meler M, King J, Leavitt S W, Lin G, Matamal R, Pendall E, Siegworf R, Van Kessel C and Ehleringer J R 2003 Tracing changes in ecosystem function under elevated carbon dioxide conditions. *BioScience* 53, 805–818.
- Peek J M, Korol J J, Gay D and Hershey T 2001 Overstory-understory biomass changes over a 35-year period in southcentral Oregon. *For. Ecol. Manag.* 150, 267–277.



- Poorter H and Navas M L 2003 Plant growth and competition at elevated CO<sub>2</sub>: on winners, losers and functional groups. *New Phytol.* 157, 175–198.
- Rainey S M, Nadelhoffer K J, Silver W L and Downs M R 1999 Effects of chronic nitrogen additions on understory species in a red pine plantation. *Ecol. Appl.* 9, 949–957.
- Reich P B 1987 Quantifying plant response to ozone: a unifying theory. *Tree Physiol.* 3, 63–91.
- Reich P B and Amundson R G 1985 Ambient levels of ozone reduce net photosynthesis in tree and crop species. *Science* 230, 566–570.
- Riegel G M, Miller R F and Krueger W C 1992 Competition for resources between understory vegetation and overstory *Pinus ponderosa* in Northeastern Oregon. *Ecol. Appl.* 2, 71–85.
- Sanders N J, Belote R T and Weltzin J F 2004 Multitropic effects of elevated atmospheric CO<sub>2</sub> on understory plant and arthropod communities. *Environ. Entomol.* 33, 1609–1616.

*Section editor: H. J. Kronzucker*