

## GAS EXCHANGE, LEAF NITROGEN, AND GROWTH EFFICIENCY OF *POPULUS TREMULOIDES* IN A CO<sub>2</sub>-ENRICHED ATMOSPHERE

PETER S. CURTIS,<sup>1,8</sup> CHRISTOPH S. VOGEL,<sup>2</sup> XIANZHONG WANG,<sup>1</sup> KURT S. PREGITZER,<sup>3</sup> DONALD R. ZAK,<sup>4</sup>  
 JOHN LUSSENHOP,<sup>5</sup> MARK KUBISKE,<sup>6</sup> AND JAMES A. TEERI<sup>7</sup>

<sup>1</sup>*Department of Evolution, Ecology, and Organismal Biology, Ohio State University, Columbus, Ohio 43210-1293 USA*

<sup>2</sup>*University of Michigan Biological Station, University of Michigan, Ann Arbor, Michigan 48109 USA*

<sup>3</sup>*School of Forestry and Wood Products, Michigan Technological University, Houghton, Michigan 49931 USA*

<sup>4</sup>*School of Natural Resources and Environment, University of Michigan, Ann Arbor, Michigan 48109 USA*

<sup>5</sup>*Department of Biology, University of Illinois, Chicago, Illinois 60680 USA*

<sup>6</sup>*School of Forestry, Mississippi State University, Mississippi State, Mississippi 39762 USA*

<sup>7</sup>*University of Michigan Biological Station, University of Michigan, Ann Arbor, Michigan 48109 USA*

**Abstract.** Predicting forest responses to rising atmospheric CO<sub>2</sub> will require an understanding of key feedbacks in the cycling of carbon and nitrogen between plants and soil microorganisms. We conducted a study for 2.5 growing seasons with *Populus tremuloides* grown under experimental atmospheric CO<sub>2</sub> and soil-N-availability treatments. Our objective was to integrate the combined influence of atmospheric CO<sub>2</sub> and soil-N availability on the flow of C and N in the plant–soil system and to relate these processes to the performance of this widespread and economically important tree species. Here we consider treatment effects on photosynthesis and canopy development and the efficiency with which this productive capacity is translated into aboveground, harvestable yield.

We grew six *P. tremuloides* genotypes at ambient (35 Pa) or elevated (70 Pa) CO<sub>2</sub> and in soil of low or high N mineralization rate at the University of Michigan Biological Station, Pellston, Michigan, USA (45°35' N, 84°42' W). In the second year of growth, net CO<sub>2</sub> assimilation rate was significantly higher in elevated-CO<sub>2</sub> compared to ambient-CO<sub>2</sub> plants in both soil-N treatments, and we found little evidence for photosynthetic acclimation to high CO<sub>2</sub>. In the third year, however, elevated-CO<sub>2</sub> plants in low-N soil had reduced photosynthetic capacity compared to ambient-CO<sub>2</sub>, low-N plants. Plants in high-N soil showed the opposite response, with elevated-CO<sub>2</sub> plants having higher photosynthetic capacity than ambient-CO<sub>2</sub> plants. Net CO<sub>2</sub> assimilation rate was linearly related to leaf N concentration (log : log scale), with identical slopes but different intercepts in the two CO<sub>2</sub> treatments, indicating differences in photosynthetic N-use efficiency. Elevated CO<sub>2</sub> increased tissue dark respiration in high-N soil (+22%) but had no significant effect in low-N soil (+9%). There were no CO<sub>2</sub> effects on stomatal conductance. At the final harvest, stem biomass and total leaf area increased significantly due to CO<sub>2</sub> enrichment in high-N but not in low-N soil. Treatment effects on wood production were largely attributable to changes in leaf area, with no significant effects on growth efficiency. We conclude that harvest intervals for *P. tremuloides* on fertile sites will shorten with rising atmospheric CO<sub>2</sub>, but that tree size at canopy closure may be unaffected.

**Key words:** carbon dioxide, elevated; feedbacks in the plant–soil system; forest responses to rising atmospheric CO<sub>2</sub>; gas exchange, plant–soil system; global climate change, ecological effects; growth efficiency; leaf nitrogen; Michigan (USA); photosynthesis; *Populus tremuloides*; respiration; soil-N availability.

### INTRODUCTION

#### *Elevated atmospheric CO<sub>2</sub> and plant–soil interactions*

The common limitation of nitrogen (N) for plant growth and carbon (C) for microbial growth in soil

results in close coupling of the plant and soil C and N cycles (Pastor and Post 1988). At a fundamental level, the response of these biogeochemical cycles to rising atmospheric CO<sub>2</sub> will be controlled by the growth response of plants and the extent to which changes in above- and belowground litter inputs alter the composition and function of microbial communities in soil. To further our understanding of plant–soil interactions with rising atmospheric CO<sub>2</sub> and hence our ability to predict changes in ecosystem function, greater attention must be paid to experiments focusing on key feedbacks

Manuscript received 16 October 1998; revised 12 April 1999; accepted 22 April 1999; final version received 20 May 1999. For reprints of this Invited Feature, see Footnote 1, p. 1.

<sup>8</sup> E-mail: curtis.7@osu.edu

in the cycling of C and N between plants and soil microorganisms (Curtis et al. 1994). These feedbacks occur within a complex chain of assimilation, transport, transformation, and release of these elements and require approaches that integrate physiological and growth processes operating in different spatial domains and time scales.

Soil microorganisms represent important sinks as well as sources of N (Vitousek and Matson 1984, Zak et al. 1990), and they are the main regulators of C dynamics within the soil (McGill et al. 1986). Increased fine-root and leaf production, and altered tissue biochemistry, could potentially alter microbial growth in soil, and thus influence the flow of N from soil microorganisms to plant roots. Although the interaction between plants and soil microorganisms has important consequences for C sequestration in terrestrial ecosystems, there are fundamental gaps in our knowledge of the mechanisms by which rising atmospheric CO<sub>2</sub> will alter the exchange of C and N between plants and soil microorganisms. For example, we understand relatively little regarding how atmospheric CO<sub>2</sub> will modify the production and timing of root C inputs to soil and how this input of substrate will influence the composition and function of microbial communities in soil. Moreover, it is likely that differences in soil-N availability will modify plant growth response to atmospheric CO<sub>2</sub>, but we have little information about this potential feedback on the flow of C and N through the soil food web. Some experimental evidence suggests that C and N cycling will slow under elevated CO<sub>2</sub> due to the production of plant litter that will enhance rates of microbial immobilization (Diaz et al. 1993, Berntson and Bazzaz 1997), whereas others have observed increased rates of C and N transformations (Zak et al. 1993, Hungate et al. 1997). Whether rising atmospheric CO<sub>2</sub> will increase, decrease, or not influence rates of C and N cycling in soil has important global implications for C sequestration in terrestrial ecosystems.

In 1993 we initiated an experiment to study the combined influence of atmospheric CO<sub>2</sub> and soil-N availability on the cycling of C and N between plant roots and soil microorganisms. We grew softwood cuttings of *Populus tremuloides* Michx. under experimental atmospheric-CO<sub>2</sub> and soil-N-availability treatments for three growing seasons in large open-top chambers. Our experiment was designed to test the broad hypothesis that there is long-term positive feedback between CO<sub>2</sub> assimilation at elevated CO<sub>2</sub>, root growth, microbial populations, and soil-N availability (Zak et al. 1993). More specifically, we hypothesized that plant C assimilation will increase in an elevated-CO<sub>2</sub> atmosphere, even under conditions of low soil-N availability. Moreover, we reasoned that this increase will be maintained over time by greater C translocation belowground and increased N acquisition. The latter will be driven by larger fine-root production and greater mycorrhizal in-

fection under CO<sub>2</sub> enrichment, resulting in greater soil exploration. Short-term soil-C availability could increase in response to greater rates of fine-root production (or turnover), and greater soil-C availability would elicit an increase in soil microbial biomass. We also predicted that net N mineralization would increase as a result of either an increase in the turnover of microbial N (via increased protozoan grazing) or through greater organic-matter mineralization by a larger microbial population.

#### *Carbon dioxide assimilation at high CO<sub>2</sub>*

Regardless of the physiological links between plants and soil microorganisms, the first stage in signal transduction from atmospheric C to biogeochemical processes in soil is the assimilation of CO<sub>2</sub> by plants. There is now a sizable body of literature detailing the gas-exchange responses of woody plants to CO<sub>2</sub> enrichment, with some convergence in estimates of the overall direction and magnitude of CO<sub>2</sub> effects on key physiological processes and allocational patterns in young (<5 yr) plants. Under optimal growth conditions (i.e., given ample water, nutrients, and light) it is clear that leaf-level light-saturated net CO<sub>2</sub> assimilation (hereafter called "A") shows a positive, sustained increase under twice-ambient CO<sub>2</sub>, with estimates from quantitative reviews ranging from +44% (Gunderson and Wullschleger 1994) to +66% (Norby et al. 1999). A sustained increase in the assimilatory capacity of leaves is central to many predictions of greater forest productivity with rising CO<sub>2</sub> (Reynolds et al. 1996). Although the effect of elevated CO<sub>2</sub> on A is probably the best understood plant CO<sub>2</sub> response at a mechanistic level, our understanding of the environmental factors causing down-regulation, or acclimation of A at high CO<sub>2</sub> ( $A_{\text{accl}}$ , sensu Gunderson and Wullschleger 1994) remains incomplete. Under some conditions root restriction elicits a strong  $A_{\text{accl}}$  response (Thomas and Strain 1991) suggesting an important role for source : sink dynamics, but this result is far from universal (McConnaughay et al. 1993). Gunderson and Wullschleger (1994) estimated a mean 21%  $A_{\text{accl}}$  across 20 studies with some indication of greater  $A_{\text{accl}}$  under low nutrients, a conclusion supported by Curtis (1996). In a more recent review, Curtis and Wang (1998) found no evidence for systematic  $A_{\text{accl}}$  by trees, except for those grown in pots <0.5 L.

A promising line of research that may improve our ability to predict  $A_{\text{accl}}$  under different environmental conditions builds on the well-recognized positive relationship between A and leaf N concentration (Field and Mooney 1986) and on the inherent tradeoffs between C and N allocation to leaf structural or defense functions on the one hand and to assimilatory capacity on the other (Lambers and Poorter 1992). The magnitude of the CO<sub>2</sub> response by A, and the probability of  $A_{\text{accl}}$ , may be largely a function of N and structural

biomass allocation within the leaf (Luo et al. 1994) and may follow predictable patterns across biomes or plant functional groups (Peterson et al. 1999). Framing the consideration of elevated-CO<sub>2</sub> effects on *A* within the context of leaf N and specific leaf area (SLA) has the additional benefit of allowing one to relate photosynthetic responses following experimental CO<sub>2</sub> manipulations to broader patterns of plant adaptation and functioning within natural ecosystems (e.g., Reich et al. 1997).

Biomass also is consistently, positively affected by elevated CO<sub>2</sub> under optimal conditions, but as growth conditions diverge from optimal our certainty in the average CO<sub>2</sub> effect size is considerably less, as evidenced by markedly different conclusions reached in recent reviews. For example, while McGuire et al. (1995) and Curtis and Wang (1998) estimated a halving of the CO<sub>2</sub> effect on biomass accumulation due to growth under low-nutrient conditions, Idso and Idso (1994) and Wullschlegel et al. (1995) concluded that there was no effect of nutrients on the magnitude of the CO<sub>2</sub> response (i.e., it was equivalent to that under optimal conditions). These different conclusions reflect both different quantitative review methods and methodological differences among the primary studies considered by the reviewers, such as the presence of interacting stress variables, length of CO<sub>2</sub> exposure, pot size, or type of exposure facility, which can affect the CO<sub>2</sub> response independently of soil nutrient status (Curtis 1996). Norby (1996) suggested that biomass gain per se may not be the best measure of woody-plant growth response to elevated CO<sub>2</sub> precisely because it is so sensitive to environmental and cultural conditions. Short-term effects on development are amplified over time in young plants so that biomass gain during early growth may be poorly related to CO<sub>2</sub> effects occurring later in stand development. As an alternative he proposed the canopy productivity index, or growth efficiency (*E*, wood production per unit leaf area per unit time) sensu Waring (1983) as a measure better tied conceptually to the control of productivity in trees. For example, as stand leaf-area index (LAI) increases, with consequent increases in competition for light, *E* typically declines in a manner reflective of species-specific adaptational characteristics and of site conditions. Factors allowing the maintenance of high *E* as LAI increases, such as shade tolerance or nutrient addition, lead to sustained increases in stand productivity during canopy closure (Waring 1983). Because CO<sub>2</sub> enrichment can affect canopy development itself (e.g., Reekie and Bazzaz 1989, Ceulemans et al. 1995, Curtis et al. 1995) *E* may be a more conservative indicator of overall performance and a better predictor of long-term behavior than is biomass gain. Indeed, among seven elevated-CO<sub>2</sub> experiments in which *E* could be calculated, the coefficient of variation (CV) for mean percentage increase in aboveground dry mass

due to elevated CO<sub>2</sub> was 75% compared to a 24% CV for the mean CO<sub>2</sub> effect on *E* (Norby 1996).

In this paper, we report the effects of elevated CO<sub>2</sub> on photosynthetic C assimilation and growth of *P. tremuloides*, the first step in the movement of C from the atmosphere to the biogeochemical cycling of C and N by soil microorganisms. Our objectives were to characterize the pattern of C gain over time as influenced by CO<sub>2</sub> and soil-N availability and to put these observations into the larger picture of tree growth dynamics and long-term stand productivity. Our null hypothesis was that C assimilation and growth would show a continued, positive response to CO<sub>2</sub> enrichment, even in low-N soil. We reasoned that greater belowground growth under elevated CO<sub>2</sub> would increase plant N acquisition, thus maintaining greater rates of C acquisition at elevated compared to ambient CO<sub>2</sub> (Zak et al. 1993).

#### METHODS

Our experiment was conducted at the University of Michigan Biological Station, Pellston, Michigan, USA, 45°35' N, 84°42' W. An array of 20 open-bottom root boxes (3.3 × 3.3 × 0.4 m) were placed in an open field in October 1993 and filled with soil. The root boxes rested on soil surface whose A and E horizons had previously been removed. Each root box was lined with 1.3-cm-thick styrofoam insulation and plastic sheeting and contained eight minirhizotron tubes (Pregitzer et al. 2000). Two soil-N-availability treatments were established by filling half of the root boxes with the A horizon of a Kalkaska series topsoil (high-N treatment), a common soil type in northern lower Michigan, and the remaining boxes with a mixture of 20% Kalkaska A horizon and 80% Rubicon C horizon sand (low-N treatment). Five centimeters of 100% Rubicon sand were placed over the soil surface of each root box to equalize surface albedo. Net nitrogen mineralization was significantly higher in the high-N soil (318 ng N·g<sup>-1</sup>·d<sup>-1</sup>) than in the low-N soil (62 ng N·g<sup>-1</sup>·d<sup>-1</sup>, Zak et al. 2000b). These N mineralization rates are well within the range normally encountered in soils of this region and where aspen would be expected to establish following disturbance (Zak et al. 2000a). Other physical and chemical properties of the two soils are presented in Table 1. Soil texture was determined using the hydrometer method. Ceramic-plate pressure membranes were used to determine soil water content at -0.03 and -1.50 MPa. Total C and N were measured using an NC2500 Elantech elemental analyzer (CE Elantech, Lakewood, New Jersey, USA). The Bray P1 method (Kuo 1996) was used to extract PO<sub>4</sub><sup>3-</sup>, and P concentrations were determined with an Alpkem RFA 300 autoanalyzer (Alpkem, Wilsonville, Oregon, USA). Soil pH was measured in a 1:1 soil-deionized water paste with a glass electrode.

Open-top chambers (3 × 2.3 m, Heagle et al. 1989)

TABLE 1. Physical and chemical properties of low- and high-N soils.

Soil properties	Low-N soil		High-N soil	
	Mean	1 SE	Mean	1 SE
Physical properties				
Sand (%)	93.5 <sup>a</sup>	0.30	72.4 <sup>b</sup>	2.14
Silt (%)	4.0 <sup>a</sup>	0.47	17.2 <sup>b</sup>	1.43
Clay (%)	2.5 <sup>a</sup>	0.50	10.1 <sup>b</sup>	1.02
Gravimetric water content, $\omega$				
-0.03 MPa	0.031 <sup>a</sup>	0.002	0.115 <sup>b</sup>	0.010
-1.50 MPa	0.017 <sup>a</sup>	0.01	0.062 <sup>b</sup>	0.005
Available water content	0.014 <sup>a</sup>	0.002	0.053 <sup>b</sup>	0.012
Chemical properties				
Total C (mg C/kg soil)	3559 <sup>a</sup>	454.7	12 489 <sup>b</sup>	2273.8
Total N (mg N/kg soil)	260 <sup>a</sup>	21.3	996 <sup>b</sup>	169.9
C:N	13.7 <sup>a</sup>	0.23	12.5 <sup>a</sup>	1.77
Extractable P (mg P/kg soil)	13.7 <sup>a</sup>	0.68	10.4 <sup>b</sup>	1.10
pH	6.74 <sup>a</sup>	0.177	6.08 <sup>b</sup>	0.127

Notes: Six soil cores (5-cm diameter) were collected to a depth of 10 cm in each open-bottom root box. Cores were composited on a chamber basis and homogenized. All analyses were conducted on the composite sample from each open-bottom root box;  $n = 10$  replicate root boxes per soil-N availability treatment. Within a row, means with the same lowercase superscript letter are not significantly different at  $P < 0.05$ .

were used to manipulate atmospheric-CO<sub>2</sub> partial pressure. Ten chambers received additional CO<sub>2</sub> (elevated-CO<sub>2</sub> treatment) and 10 chambers received no additional CO<sub>2</sub> (ambient-CO<sub>2</sub> treatment). Carbon dioxide treatments were randomized across soil-N availability treatments within five replicate blocks. Carbon dioxide partial pressure was increased by dispensing 100% CO<sub>2</sub> into an input blower via manual flowmeters with the atmosphere inside all elevated-CO<sub>2</sub> chambers and one ambient-CO<sub>2</sub> chamber monitored continuously using an infrared gas analyzer. Carbon dioxide treatments were maintained 24 hr/d for all days during which green leaves were present in the chambers, but CO<sub>2</sub> fumigation was terminated following leaf senescence in 1994 and 1995. Daytime (0701–1900) elevated-CO<sub>2</sub> treatment was  $70.7 \pm 0.06$  Pa (mean of daily averages across all chambers and days  $\pm 1$  SE) and nighttime (1901–0700) elevated-CO<sub>2</sub> treatment was  $73.2 \pm 0.14$  Pa. The ambient-CO<sub>2</sub> treatment was  $35.7 \pm 0.06$  Pa (day) and  $38.32 \pm 0.09$  Pa (night). Light and temperature were continuously monitored every 10 min inside and outside chambers during the growing season. Daytime temperatures were  $1.34 \pm 0.24^\circ\text{C}$  (mean  $\pm 1$  SE) higher inside than outside chambers across the three growing seasons. The chamber plastic transmitted 81% of ambient PAR (photosynthetically active radiation).

On 7 June 1994 two individuals from each of six locally derived trembling aspen (*Populus tremuloides* Michx.) genotypes were transplanted into each root box (=12 trees per chamber) and CO<sub>2</sub> treatments begun. Genotype was therefore a subplot within the CO<sub>2</sub>  $\times$  soil-N whole plot, yielding a split-plot randomized complete-block design. The genotypes were selected based on field observations of patterns of autumnal leaf senescence. Three genotypes (42E, 51E, and 61E) typically dropped their leaves in mid-October (early leaf-

drop phenotype) and three genotypes (1L, 2L, and 8L) typically dropped their leaves in early November (late leaf-drop phenotype). By selecting these genotypes we hoped to encompass the variation present locally in a phenological trait that might influence the CO<sub>2</sub> response.

Individuals were arranged in two concentric circles within each chamber—an outer circle of eight trees and an inner circle of four trees. During June and July 1994 chambers were irrigated daily with 7–10 L water/m<sup>2</sup> soil surface, except on rainy or cloudy, cool days. Plants relied on natural precipitation from August 1994 until the beginning of the harvest in July 1996. In September 1995 all chambers were enlarged to 3.4 m in height to accommodate tree growth. During the following winter 5% of each tree's total aboveground wood volume was removed by pruning a portion of the terminal shoot. This was done to ensure containment of all trees within the open-top chambers until the end of the experiment.

#### Gas exchange

Leaf net CO<sub>2</sub> assimilation and stomatal conductance ( $g_s$ ) were measured on 12 August 1995, and on 6 June and 16 July 1996 using a LI-COR LI-6400 portable photosynthesis system (LI-COR, Lincoln, Nebraska). Leaf temperature was maintained at 26°–30°C and saturating light intensity (1800–2000  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  PAR) supplied with a cuvette illuminator. Gas-exchange measurements were performed between 1000 and 1600 on young, fully expanded leaves that had developed in full sun. CO<sub>2</sub> enriched air was supplied to the LI-6400 for measurements above ambient-CO<sub>2</sub> partial pressure ( $p(\text{CO}_2)$ ).

Both  $A$  and  $g_s$  were measured at growth  $p(\text{CO}_2)$  and at internal leaf  $p(\text{CO}_2)$  ( $C_i$ ) of 27 and 56 Pa. These  $C_i$

values were representative of full-sun leaves from ambient- and elevated-CO<sub>2</sub> treatments, respectively, and allowed us to evaluate photosynthetic capacity independent of environmental effects on  $g_s$ . Leaves were first exposed to their growth p(CO<sub>2</sub>) until steady-state gas exchange was observed, then measurements made at 27 Pa C<sub>r</sub>, followed by 56 Pa C<sub>r</sub>. At each measurement p(CO<sub>2</sub>), leaves were allowed to equilibrate for at least 4 min before measurements of  $A$  (leaf-level light-saturated net CO<sub>2</sub> assimilation) or  $g_s$  (stomatal conductance) were made. Following each measurement, 2.6-cm<sup>2</sup> leaf discs were collected from the sampled leaf, frozen on dry ice, and later lyophilized to dryness. These samples were used for calculation of SLA (specific leaf area) and of leaf N on a mass (N<sub>m</sub>) or an area (N<sub>a</sub>) basis. Tissue N was measured with a CE Elantech CN 1200 elemental analyzer. Leaf discs were also collected from mature, high-light-grown leaves, embedded in paraffin, and 10- $\mu$ m sections examined using light microscopy for determination of leaf thickness.

Leaf respiration was measured on shaded leaves during the day once in 1995 (3 July, blocks 2–5), twice in 1996 (5 June and 9 July, blocks 2–5), and once on shaded leaves at night in 1996 (5 June, blocks 1–5). All leaf-respiration measurements were made at 28°C and at growth p(CO<sub>2</sub>) using the LI-6400. In 1995 only genotypes 1L, 51E, and 61E were measured while in 1996 all genotypes were measured. Individuals were selected at random within a chamber, resulting in a partial repeated measure of individuals although branches sampled were always different. Stem respiration was measured once in 1996 (30 June–5 July, blocks 3–5, genotypes 1L and 61E) using the LI-6400 fitted with a custom branch cuvette; surface temperatures varied between 26° and 30°C and measurements were made at growth p(CO<sub>2</sub>).

#### Growth measurements

In 1994 the total leaf area per plant (LA) was calculated by summing the area of individual leaves (LA<sub>*i*</sub>), which was estimated by

$$LA_i = x_i \times (L_i \times L_w) + y_i \quad (1)$$

where  $L_i$  was leaf length,  $L_w$  was leaf width, and  $x_i$  and  $y_i$  were genotype-specific regression coefficients obtained from destructive harvest of non-experimental plants. In 1995 LA was calculated by summing the leaf area of individual branches (LA<sub>*b*</sub>), estimated by

$$LA_b = w_b \times B_d + x_b \times B_l + y_b \times (B_l \times B_d) + z_b, \quad (2)$$

where  $B_l$  was branch length,  $B_d$  was branch basal diameter, and  $w_b$ ,  $x_b$ ,  $y_b$ , and  $z_b$  were genotype-specific regression coefficients obtained from measuring  $L_i$  and  $L_w$  on all leaves from a subset of experimental branches and applying Eq. 1. Stem volume was estimated from  $B_l$  and  $B_d$  assuming a conical stem geometry.

Growth efficiency was calculated both on the basis

of estimated yearly stem-volume increase (above-ground volume growth efficiency,  $E_v$ ) and on the basis of harvested stem dry mass ( $E_{dm}$ ). All growth-efficiency calculations were made on the basis of growth per chamber (as opposed to growth per plant) of the four plants growing in the center of the chamber, thereby eliminating the influence of side light, using the following expressions:

$$E_v = \frac{V_n - V_{n-1}}{d_n \times LA_n} \quad (3)$$

$$E_{dm} = \frac{S_{dm}^{1996}}{401 \times \sum_{n=1994} LA_n} \quad (4)$$

where  $V_n$  was estimated stem volume in year  $n$ ,  $LA_n$  was estimated leaf area in year  $n$ ,  $d_n$  was number of days since leaf out in year  $n$ , and  $S_{dm}$  was harvested stem dry mass. There were 401 d of leaf exposure during the entire study ( $=\sum d_n$ ).

Plants were destructively harvested beginning 8 July 1996. Leaves were separated into three canopy-position classes based on their height above the soil surface: <1 m, 1–2 m, and >2 m for the high-N treatment, and <0.5 m, 0.5–1 m, and >1 m for the low-N treatment. Total leaf fresh mass per tree was obtained for each position class, the leaves were composited, and a subsample taken for leaf-area measurement (LI-COR LI-3000 leaf-area meter). This subsample was dried at 65°C and used for fresh mass : dry mass and mass : leaf area conversions for each tree. Leaf tissue C and N concentrations were determined from these subsamples using a CE Elantech CN 1200 elemental analyzer. Stem tissue was separated at each year's terminal bud scale scar, yielding first-year wood (produced in 1994, secondary growth in 1995 and 1996), second-year wood (produced in 1995, secondary growth in 1996), and third-year wood (produced in 1996). One ~10-cm subsample was cut from each age class and air dried for density determination. Stem volume at harvest for each individual was determined from stem age-class dry mass and stem age-class density. Zak et al. (2000a) and Pregitzer et al. (2000) provide details regarding the belowground harvest of this experiment.

#### Carbohydrate analysis

Leaf discs (2.6 cm<sup>2</sup>) were excised between 1400 and 1500 on 3 July 1996 from the youngest fully expanded leaf on a branch exposed to full sun, immediately frozen in liquid nitrogen, and stored at –80°C until lyophilized. Powdered, lyophilized samples were extracted with 80% ethanol at 80°C, the supernatant evaporated to dryness, and then redissolved in H<sub>2</sub>O + polyvinylpyrrolidone. Soluble carbohydrates were analyzed enzymatically using a modification of the procedure of Jones et al. (1977). The recovery rate was 95%. The ethanol-extracted tissue pellet was suspended

TABLE 2. Gas exchange of *Populus tremuloides* grown in low- or high-N soil and at ambient or elevated CO<sub>2</sub>.

Date	Parameter	Low-N soil				High-N soil			
		Ambient CO <sub>2</sub>		Elevated CO <sub>2</sub>		Ambient CO <sub>2</sub>		Elevated CO <sub>2</sub>	
		Mean	1 SE	Mean	1 SE	Mean	1 SE	Mean	1 SE
August 1995	A <sub>gr</sub>	21.0 <sup>c</sup>	1.4	26.4 <sup>ab</sup>	1.1	20.9 <sup>bc</sup>	1.9	32.6 <sup>a</sup>	2.1
	g <sub>sgr</sub>	0.64 <sup>a</sup>	0.07	0.44 <sup>a</sup>	0.06	0.55 <sup>a</sup>	0.11	0.46 <sup>a</sup>	0.06
	A <sub>27</sub>	21.5 <sup>ab</sup>	1.1	17.5 <sup>c</sup>	0.35	23.4 <sup>a</sup>	0.6	20.2 <sup>bc</sup>	1.0
	g <sub>s27</sub>	0.65 <sup>a</sup>	0.06	0.41 <sup>a</sup>	0.04	0.54 <sup>a</sup>	0.12	0.42 <sup>a</sup>	0.07
	A <sub>56</sub>	30.8 <sup>ab</sup>	1.2	26.5 <sup>b</sup>	1.1	34.8 <sup>a</sup>	0.9	33.7 <sup>a</sup>	2.3
	g <sub>s56</sub>	0.54 <sup>a</sup>	0.06	0.43 <sup>a</sup>	0.05	0.49 <sup>a</sup>	0.12	0.46 <sup>a</sup>	0.08
July 1996	A <sub>gr</sub>	9.1 <sup>b</sup>	1.0	12.1 <sup>b</sup>	1.6	9.2 <sup>b</sup>	0.8	21.0 <sup>a</sup>	1.9
	g <sub>sgr</sub>	0.15 <sup>a</sup>	0.04	0.11 <sup>a</sup>	0.03	0.13 <sup>a</sup>	0.02	0.16 <sup>a</sup>	0.04
	A <sub>27</sub>	11.9 <sup>a</sup>	0.8	6.7 <sup>b</sup>	0.7	12.9 <sup>a</sup>	1.5	15.1 <sup>a</sup>	1.2
	g <sub>s27</sub>	0.15 <sup>a</sup>	0.04	0.10 <sup>a</sup>	0.02	0.13 <sup>a</sup>	0.02	0.13 <sup>a</sup>	0.04
	A <sub>56</sub>	18.2 <sup>b</sup>	0.6	13.7 <sup>c</sup>	1.0	20.5 <sup>b</sup>	1.1	24.6 <sup>a</sup>	1.3
	g <sub>s56</sub>	0.14 <sup>a</sup>	0.03	0.11 <sup>a</sup>	0.03	0.11 <sup>a</sup>	0.02	0.15 <sup>a</sup>	0.04

Notes: Measurements of net CO<sub>2</sub> assimilation (A) and stomatal conductance (g<sub>s</sub>) were made at growth p(CO<sub>2</sub>) (A<sub>gr</sub>, g<sub>sgr</sub>) and at internal p(CO<sub>2</sub>) of 27 Pa (A<sub>27</sub>, g<sub>s27</sub>) and 56 (A<sub>56</sub>, g<sub>s56</sub>) Pa, respectively [p(CO<sub>2</sub>) partial pressure]; n = 5 replicate chambers per CO<sub>2</sub> and soil-N availability treatment. Within a row, means with the same lowercase superscript letters are not significantly different at P < 0.05.

in 0.2 mol/L KOH, boiled for 20 min, and brought to pH 7.0 with 1.0 mol/L CH<sub>3</sub>COOH. Amyloglucosidase (EC [Enzyme Classification] 3.2.1.3) was added to the resuspended pellet and incubated for 1 h at 55°C. Starch concentration was determined as glucose equivalents using the same procedure as for soluble sugars. Starch recovery was 93%.

#### Statistical analysis

Data were analyzed by analysis of variance (ANOVA) for a split-plot factorial block design where main effects (CO<sub>2</sub>, and soil fertility) and CO<sub>2</sub> × soil fertility interaction were tested over the CO<sub>2</sub> × fertility nested within block mean square. The effect of genotype and all treatment interactions with genotype were tested over the error mean square. Growth efficiency, which was expressed on a per chamber basis, was analyzed as a two-way factorial ANOVA with five replicate blocks. Comparison among CO<sub>2</sub> and fertility treatment means was by least significant difference for a priori comparisons (between CO<sub>2</sub> treatments within a fertility treatment), and by minimum significant difference for all a posteriori comparisons (Sokal and Rohlf 1981).

The overall effect of CO<sub>2</sub> on respiration across years and tissue types was evaluated by calculating the 95% confidence interval around the mean CO<sub>2</sub> effect ( $\bar{L}$ ), where

$$\bar{L} = \frac{\sum_{i=1}^k w_i L_i}{\sum_{i=1}^k w_i} \quad (5)$$

and  $L_i = \log(\bar{X}_{ei}/\bar{X}_{ai})$ , the log-transformed ratio of elevated ( $\bar{X}_{ei}$ ) to ambient ( $\bar{X}_{ai}$ ) CO<sub>2</sub> treatment means of the *i*th set of measurements, and  $w_i$  is the reciprocal of

the total variance of  $L_i$  (Hedges et al. 1999). This is, therefore, a mean estimate weighted according to the precision (standard error) of the *k* separate studies and is typical for meta-analyses of this kind (Cooper and Hedges 1994).

The effect of CO<sub>2</sub> enrichment on the relationship between A and leaf N was evaluated using the linear model:

$$\log(A)_i = \beta_0 + \beta_1 \log(\text{leaf N})_i + \beta_2 X_i + \beta_3 \log(\text{leaf N})_i X_i + \varepsilon_i \quad (6)$$

where  $\beta_0$  is the centered y intercept for ambient CO<sub>2</sub>,  $\beta_1$  is the slope for the ambient-CO<sub>2</sub> treatment,  $\beta_2$  and  $\beta_3$  are the change in the centered y intercept and slope, respectively, due to CO<sub>2</sub> enrichment,  $X_i$  is a dummy variable coded 0 for ambient CO<sub>2</sub> and 1 for elevated CO<sub>2</sub>, and  $\varepsilon_i$  is residual error (Peterson et al. 1999). Significance of coefficients was determined by least-squares regression.

## RESULTS

### Gas exchange

Net CO<sub>2</sub> assimilation varied both among treatments and across years (Table 2). In mid-August 1995, A at growth p(CO<sub>2</sub>) (A<sub>gr</sub>) increased significantly in both soil N treatments (+56% and +26% in high- and low-N soil, respectively), but there was no effect of soil-N availability within CO<sub>2</sub> treatments. High-CO<sub>2</sub>-grown plants in both soil N treatments had reduced photosynthetic capacity at 27 Pa C<sub>i</sub> relative to ambient-CO<sub>2</sub>-grown plants, but no reduction in capacity at 56 Pa C<sub>i</sub>. There was no CO<sub>2</sub> or soil-N availability effect on g<sub>s</sub> under any measurement C<sub>i</sub> (Table 2).

By mid-July 1996 this picture had changed substantially (Table 2). Both A and g<sub>s</sub> were significantly lower

TABLE 3. Analysis of variance of leaf area and stem dry mass ( $S_{dm}$ ) per plant at the final harvest and for net CO<sub>2</sub> assimilation at growth p(CO<sub>2</sub>) ( $A_{gr}$ ) measured in July 1996.

Source of variation	df	Leaf area		$S_{dm}$		$A_{gr}$	
		MS	$F$	MS	$F$	MS	$F$
CO <sub>2</sub>	1	5.8	10.4**	317 969	28.3***	1320	50.8***
Soil N	1	165.0	297.5***	7 437 212	663.2***	490	18.9**
CO <sub>2</sub> × Soil N	1	5.9	10.6**	219 824	19.6***	461	17.7**
Block	4	0.7	0.9	69 538	2.5	111	7.1***
Block (CO <sub>2</sub> × Soil N)	12	0.6		11 215		26	
Genotype	5	6.6	8.0***	324 596	11.6***	20	1.3
CO <sub>2</sub> × Genotype	5	1.1	1.3	19 532	0.7	7.2	0.5
Genotype × Soil N	5	4.3	5.3***	194 822	6.9***	9.0	0.5
CO <sub>2</sub> × Soil N × Genotype	5	0.5	0.6	19 732	0.7	5.9	0.4
Error	80	0.6		28 062		15.7	

Notes: Sources of variation were CO<sub>2</sub> treatment (ambient or elevated), soil-N availability (low or high), and genotype (six genotypes total). For  $A_{gr}$ , block df = 3.

\*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ .

in 1996 than in 1995, and the effect of CO<sub>2</sub> enrichment on  $A$  had diverged between the two soil-N-availability treatments. In high-N soil,  $A_{gr}$  was 128% greater in elevated-compared to ambient-CO<sub>2</sub>-grown plants, while there was no significant CO<sub>2</sub> effect in low-N soil. This CO<sub>2</sub> × soil-N interaction also was reflected in the measurements of  $A$  at constant  $C_i$ . In low-N soil, elevated-CO<sub>2</sub>-grown plants had significantly lower  $A$  at both measurement  $C_i$ 's than did ambient-CO<sub>2</sub>-grown plants. In high-N soil, however,  $A$  in elevated-CO<sub>2</sub>-grown plants was equal to that in ambient-CO<sub>2</sub>-grown plants when measured at 27 Pa CO<sub>2</sub>, and greater than that in ambient-CO<sub>2</sub>-grown plants when measured at 56 Pa CO<sub>2</sub>. As in 1995, there were no treatment effects on  $g_s$ . In July 1996 we found no evidence for differential photosynthetic capacity among genotypes or in genotypic responses to the CO<sub>2</sub> or soil N treatments (Table 3).

Treatment effects on dark respiration were somewhat variable, with their magnitude changing over time, and between leaf and stem tissue (Table 4). In general, elevated-CO<sub>2</sub>-grown plants had higher respiration rates

than did ambient-CO<sub>2</sub>-grown plants, the single exception being for stem respiration in plants from low-N soil. Because of low sample sizes, however, none of the individual CO<sub>2</sub> responses was statistically significant. To gain an estimate of the overall effect of CO<sub>2</sub> on aboveground respiration we combined these individual data sets, weighting each by their precision. The resulting mean CO<sub>2</sub> effect size suggests a greater positive respiratory response to CO<sub>2</sub> in high-N soil than in low-N soil, the latter being not significantly different from zero (95% CI overlaps zero). These confidence limits have some uncertainty associated with them since there was at least partial non-independence among individual data sets. However, applying a more conservative Type I error probability of 1% does not alter the result (CI of -2% to +22% for the low-N mean and 5% to 34% for the high-N mean).

Net assimilation in 1996, expressed on a tissue mass basis, showed a strong positive relationship with  $N_m$  (Fig 1). Plotted on a log : log scale this relationship was linear, with elevated- and ambient-CO<sub>2</sub> treatments having identical slopes of 1.39 at growth p(CO<sub>2</sub>) and 1.31

TABLE 4. Dark respiration in leaves and stems of *Populus tremuloides* grown in low- or high-N soil and at ambient or elevated CO<sub>2</sub>.

Tissue sample	Low-N soil			High-N soil		
	Ambient CO <sub>2</sub> ( $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )	Elevated CO <sub>2</sub> ( $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )	Change (%)†	Ambient CO <sub>2</sub> ( $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )	Elevated CO <sub>2</sub> ( $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )	Change (%)†
Leaf, 1995 day	2.49 (0.22)	3.12 (0.12)	+25.3	3.02 (0.28)	3.48 (0.27)	+15.2
Leaf, 1996a day	2.62 (0.17)	2.94 (0.20)	+12.2	2.67 (0.13)	3.10 (0.24)	+16.1
Leaf, 1996b day	1.96 (0.06)	2.10 (0.14)	+7.1	1.77 (0.13)	2.09 (0.23)	+18.1
Leaf, 1996 night	1.68 (0.13)	1.69 (0.10)	+5.9	1.47 (0.17)	1.95 (0.31)	+32.6
Stem, 1996 day	4.53 (0.74)	3.82 (0.48)	-15.7	3.50 (0.50)	5.40 (0.41)	+54.3
$\bar{L}$ (%)‡			+9.0			+21.8
(95% CI)			(0-19)			(9-36)

Notes: Leaves were measured once in 1995 during the day, twice in 1996 during the day (1996a and 1996b), and once in 1996 during the night; stems were measured once in 1996 during the day. Respiration data are means with 1 SE in parentheses;  $n = 3-5$  replicate chambers per CO<sub>2</sub> and soil-N availability treatment.

† Percentage change due to CO<sub>2</sub> treatment.

‡ The mean CO<sub>2</sub> effect size,  $\bar{L}$ , was calculated from the log-transformed ratios of elevated to ambient means within a soil N treatment (see *Methods: Statistical analysis* for details).




FIG. 1. Change in net CO<sub>2</sub> assimilation in relation to leaf N of *Populus tremuloides* growing in low-N (open symbols) or high-N (solid symbols) soil and at ambient (circles, dashed line) or elevated (squares, solid line) CO<sub>2</sub>. Note log scales. Measurements of A (leaf-level light-saturated net CO<sub>2</sub> assimilation) were made in 1996 on day of year 157 (symbols without +) or day of year 197 (symbols with +) and both at growth p(CO<sub>2</sub>) (A) and 56 Pa C<sub>i</sub> (B). Each point represents the mean response of one chamber with lines fitted by least-squares regression.

at 56 Pa C<sub>i</sub>. There was a significant difference in intercept for lines fitted to elevated- vs. ambient-CO<sub>2</sub> plants at growth p(CO<sub>2</sub>) (Fig. 1A) but not at a common C<sub>i</sub> of 56 Pa (Fig. 1B) or 27 Pa (data not shown). Note that tissue N concentration and A declined in all treatments between the day of year (DOY) 157 and DOY197 sampling dates.

#### Leaf tissue chemistry

Leaf N showed a consistent response to the treatments across the latter two years of the experiment (Table 5). There was a reduction in N<sub>m</sub> (leaf N on a mass basis) due both to CO<sub>2</sub> enrichment and low soil-N availability, leading to increases in tissue C/N. For N<sub>a</sub> (leaf N on an area basis), however, the relative effect of CO<sub>2</sub> was substantially less, becoming significant only in low-N soil. Leaf total non-structural carbohydrate (TNC) increased with elevated CO<sub>2</sub> and low-N soil, due primarily to increased starch concentration (Fig. 2). This increase, together with parallel treatment effects on leaf thickness led to significant reductions in specific leaf area (SLA) in elevated compared with ambient-CO<sub>2</sub>-grown plants (Table 5). Changes in leaf TNC accounted for ~58% of the decrease in SLA in low-N soil and ~54% of the decrease in high-N soil.

#### Aboveground growth

Aboveground growth was modest in the first year of the experiment, as plants acclimated to the chamber environment. During the first year of growth, leaf area and stem volume increased in high-N soil compared to low-N soil but showed no response to CO<sub>2</sub> enrichment (Fig. 3, Table 6). During the second year, however, aboveground growth was dramatic, particularly in high-N soil where stem volume increased over 50-fold

TABLE 5. Leaf chemical and physical properties of *Populus tremuloides* grown in low- or high-N soil and at ambient or elevated CO<sub>2</sub>.

Leaf property	Low-N soil				High-N soil			
	Ambient CO <sub>2</sub>		Elevated CO <sub>2</sub>		Ambient CO <sub>2</sub>		Elevated CO <sub>2</sub>	
	Mean	1 SE	Mean	1 SE	Mean	1 SE	Mean	1 SE
August 1995								
N <sub>m</sub> (%)	2.1 <sup>a</sup>	0.1	1.6 <sup>c</sup>	0.1	2.7 <sup>b</sup>	0.1	2.1 <sup>a</sup>	0.1
N <sub>a</sub> (g/m <sup>2</sup> )	1.3 <sup>c</sup>	0.1	1.1 <sup>b</sup>	0.03	1.8 <sup>a</sup>	0.1	1.7 <sup>a</sup>	0.1
C (%)	50.5 <sup>ac</sup>	0.1	50.1 <sup>c</sup>	0.2	51.0 <sup>bd</sup>	0.1	50.3 <sup>acc</sup>	0.1
C/N	25.1 <sup>a</sup>	1.3	33.3 <sup>c</sup>	1.7	19.4 <sup>b</sup>	0.4	24.7 <sup>a</sup>	0.6
SLA (m <sup>2</sup> /kg)	15.8 <sup>d</sup>	0.4	13.8 <sup>ce</sup>	0.7	14.9 <sup>bcd</sup>	0.3	12.0 <sup>ac</sup>	0.6
July 1996								
N <sub>m</sub> (%)	1.8 <sup>d</sup>	0.1	1.4 <sup>c</sup>	0.1	2.5 <sup>b</sup>	0.1	2.2 <sup>a</sup>	0.04
N <sub>a</sub> (g/m <sup>2</sup> )	1.3 <sup>c</sup>	0.1	1.2 <sup>b</sup>	0.03	1.7 <sup>a</sup>	0.04	1.7 <sup>a</sup>	0.04
C (%)	50.8 <sup>a</sup>	0.1	50.6 <sup>a</sup>	0.3	50.8 <sup>a</sup>	0.3	50.4 <sup>a</sup>	0.1
C/N	29.3 <sup>c</sup>	1.3	38.3 <sup>b</sup>	1.7	20.6 <sup>a</sup>	0.8	23.8 <sup>a</sup>	0.6
SLA (m <sup>2</sup> /kg)	13.5 <sup>de</sup>	0.1	12.1 <sup>c</sup>	0.3	14.6 <sup>be</sup>	0.2	12.8 <sup>acd</sup>	0.4
L <sub>t</sub> (μm)	136 <sup>ad</sup>	3.1	146 <sup>b</sup>	1.7	128 <sup>cd</sup>	3.7	137 <sup>ab</sup>	2.9

Notes: Leaves were sampled for N, C, and specific leaf area (SLA) following gas-exchange measurements in 1995 and 1996 and for leaf thickness (L<sub>t</sub>) prior to harvest in 1996. Leaf N is expressed on both a mass (N<sub>m</sub>) and an area (N<sub>a</sub>) basis; n = 5 replicate chambers per CO<sub>2</sub> and soil-N availability treatment. Within a row means with the same lowercase superscript letter are not significantly different at P < 0.05.






FIG. 2. The concentration of starch, soluble sugar (Sol. Sugar), and total non-structural carbohydrates (TNC) in *Populus tremuloides* leaves after 2.5 seasons of growth in low-N (open bars) or high-N (solid bars) soil and at ambient (A) or elevated (E) CO<sub>2</sub>. Data are means and 1 SE;  $n = 5$  replicate chambers per CO<sub>2</sub> and soil-N-availability treatment.

compared to a ~7-fold increase in low-N soil. Carbon dioxide enrichment resulted in a significant increase in both leaf area and stem volume in year 2, but only in high-N soil. In that soil treatment, the increase in stem volume was nearly equally distributed between second-year growth (+28%) and first-year growth (+35%). Second-year stem volume was over twice that of first-year stem volume in low-N soil but only 20–30% greater in high-N soil.

The CO<sub>2</sub> and soil-N-availability effects on growth observed in year two continued into year three (Fig. 3, Table 6). In high-N soil, elevated CO<sub>2</sub> increased both leaf area (+28%) and stem volume (+37%), but CO<sub>2</sub> had no effect in low-N soil. In high-N soil, all yearly growth classes showed increased stem volume at elevated compared to ambient CO<sub>2</sub>. Wood density increased significantly in low- compared to high-N soil and due to CO<sub>2</sub> enrichment in first- and second-year wood in low-N soil (Table 6). Elevated CO<sub>2</sub> caused an increase in harvested stem mass in all yearly growth classes in high-N soil, resulting in a 34% increase in total aboveground wood production. In low-N soil, no year class responded significantly to CO<sub>2</sub> enrichment, and total wood production was consequently unaffected by CO<sub>2</sub> treatment.

This pattern of treatment response was reflected in the ANOVA of harvested leaf area and stem mass (Table 3). The effects of CO<sub>2</sub> and soil-N availability were highly significant, as was the interaction between atmospheric CO<sub>2</sub> and soil-N availability. The effect of genotype also was significant for these measures, as was the genotype  $\times$  soil N interaction. For example, in low-N soil there was no difference in final leaf area among genotypes while in high-N soil genotype 1L had over twice the leaf area as genotypes 42E and 51E (Fig. 4). A similar response was observed for final stem vol-

ume and mass, and for stem volume and leaf area in year two (data not shown). At the final harvest, we found no genotype  $\times$  CO<sub>2</sub> or genotype  $\times$  CO<sub>2</sub>  $\times$  soil N interaction, indicating a similar aboveground growth response to CO<sub>2</sub> by all genotypes in each soil N treatment.

The vertical distribution of leaf area in year three was not markedly affected by CO<sub>2</sub> treatment (Fig. 5). In high-N soil, CO<sub>2</sub> enrichment increased leaf area throughout the canopy, with a somewhat greater response in the upper third of the canopy compared with the lower sections. However, the relative distribution of leaf area among canopy-position classes was unchanged. There was no effect of CO<sub>2</sub> on leaf area in any canopy position in low-N soil.

Stem volume growth efficiency increased with increasing chamber leaf-area index (LAI) across the three years of the experiment (Fig. 6). Although final LAI was highest at elevated CO<sub>2</sub> and in high-N soil, and trees in high-N soil had over twice the leaf area as did those in low-N soil, we were unable to distinguish separate functional relationships between  $E_v$  (aboveground volume growth efficiency) and LAI for the different treatment combinations. That is, all treatments appeared to have similar  $E_v$  at equivalent LAI. Stem mass growth efficiency, obtained at the final harvest, showed




FIG. 3. Change in leaf area (A) and stem volume (B) of *Populus tremuloides* during 2.5 seasons of growth in low-N (open symbols) or high-N (solid symbols) soil and at ambient (circles) or elevated (squares) CO<sub>2</sub>. (In some cases open circles are completely hidden from view by open squares.) Data are means  $\pm$  1 SE;  $n = 5$  replicate chambers per CO<sub>2</sub> and soil-N-availability treatment.

TABLE 6. Stem volume ( $S_v$ ), stem density ( $S_d$ ), and stem dry mass ( $S_{dm}$ ) per tree of *Populus tremuloides* grown in low- or high-N soil and at ambient or elevated  $CO_2$ .

Year	Stem parameter	Low-N soil				High-N soil			
		Ambient $CO_2$		Elevated $CO_2$		Ambient $CO_2$		Elevated $CO_2$	
		Mean	1 SE	Mean	1 SE	Mean	1 SE	Mean	1 SE
1994	Total $S_v$ (cm <sup>3</sup> )	1.4 <sup>b</sup>	0.2	1.8 <sup>b</sup>	0.6	12.8 <sup>a</sup>	3.9	16.2 <sup>a</sup>	3.0
1995	$S_{v1}$ (cm <sup>3</sup> )	35.7 <sup>b</sup>	4.4	41.4 <sup>b</sup>	9.7	318.6 <sup>a</sup>	42.4	431.0 <sup>a</sup>	62.0
	$S_{v2}$ (cm <sup>3</sup> )	83.7 <sup>c</sup>	8.9	84.3 <sup>c</sup>	17.3	411.5 <sup>b</sup>	29.3	527.5 <sup>a</sup>	41.4
	Total $S_v$ (cm <sup>3</sup> )	119.4 <sup>c</sup>	13.1	125.7 <sup>c</sup>	25.3	730.1 <sup>b</sup>	52.2	958.5 <sup>a</sup>	99.4
1996	$S_{v1}$ (cm <sup>3</sup> )	81.7 <sup>c</sup>	10.0	100.3 <sup>c</sup>	15.1	480.3 <sup>b</sup>	56.5	648.3 <sup>a</sup>	77.2
	$S_{v2}$ (cm <sup>3</sup> )	205.6 <sup>c</sup>	19.3	204.3 <sup>c</sup>	33.4	829.4 <sup>b</sup>	28.2	1094.9 <sup>a</sup>	44.2
	$S_{v3}$ (cm <sup>3</sup> )	82.2 <sup>c</sup>	10.6	92.2 <sup>c</sup>	21.5	265.7 <sup>b</sup>	22.8	400.2 <sup>a</sup>	32.3
	Total $S_v$ (cm <sup>3</sup> )	370.8 <sup>c</sup>	34.3	396.3 <sup>c</sup>	65.7	1575.4 <sup>b</sup>	70.3	2150.2 <sup>a</sup>	109.4
	$S_{d1}$ (g/cm <sup>3</sup> )	0.444 <sup>c</sup>	0.010	0.472 <sup>b</sup>	0.011	0.388 <sup>a</sup>	0.005	0.395 <sup>a</sup>	0.002
	$S_{d2}$ (g/cm <sup>3</sup> )	0.400 <sup>c</sup>	0.007	0.449 <sup>b</sup>	0.010	0.361 <sup>a</sup>	0.007	0.338 <sup>a</sup>	0.006
	$S_{d3}$ (g/cm <sup>3</sup> )	0.409 <sup>a</sup>	0.012	0.502 <sup>a</sup>	0.043	0.367 <sup>a</sup>	0.008	0.414 <sup>a</sup>	0.059
	$S_{dm1}$ (g)	35.0 <sup>c</sup>	4.2	43.9 <sup>c</sup>	6.1	182.1 <sup>b</sup>	19.2	250.6 <sup>a</sup>	31.2
	$S_{dm2}$ (g)	78.6 <sup>c</sup>	8.1	82.6 <sup>c</sup>	11.1	282.8 <sup>b</sup>	5.6	356.4 <sup>a</sup>	14.2
	$S_{dm3}$ (g)	32.0 <sup>c</sup>	3.6	37.5 <sup>c</sup>	7.7	93.0 <sup>b</sup>	7.8	139.4 <sup>a</sup>	9.3
Total $S_{dm}$ (g)	145.6 <sup>c</sup>	13.9	164.1 <sup>c</sup>	22.2	557.9 <sup>b</sup>	22.4	746.5 <sup>a</sup>	47.7	

Notes: In 1994 and 1995,  $S_v$  was estimated from non-destructive branch measurements and in 1996 from harvest data. In 1995 and 1996, measurements were further subdivided according to the year class of growth (subscript 1 = year class 1, etc.); n = 5 replicate chambers per  $CO_2$  and soil-N availability treatment. Within a row, means with the same lowercase superscript letter are not significantly different at  $P < 0.05$ .

a similar pattern, with no significant  $CO_2$  effects on  $E_{dm}$  within soil N treatments (Fig. 6). Similar results were obtained when all trees, not just the central individuals in each chamber, were included in the analysis (data not shown).

#### DISCUSSION

Elevated- $CO_2$  effects on  $A$  (leaf-level light-saturated net  $CO_2$  assimilation) in our study followed a pattern





FIG. 4. Genotypic differences in leaf-area growth in soils of differing N availability in *Populus tremuloides*. Genotypes are distinguished by different symbols and whether they are of early (open symbols) or late (solid symbols) leaf-drop phenotype. Vertical bars indicate groups of means that are not significantly different from one another at  $P < 0.05$ .

typical of many woody species, with multi-year increases in  $A_{gr}$  ( $A$  at growth  $CO_2$  partial pressure) (Norby et al. 1992, Tissue et al. 1993, Eamus et al. 1995) but also variation in the magnitude of this response temporally (Gunderson et al. 1993, Lewis et al. 1996) and with soil fertility (Tissue et al. 1993). In high-N soil, we found little evidence for  $A_{accl}$  (acclimation of  $A$  at high  $CO_2$ ) in elevated- $CO_2$ -grown plants, similar to the results of Curtis et al. (1995), Will and Ceulemans (1997), and Kalina and Ceulemans (1997), all working with hybrid poplar. In low-N soil, however,  $A_{accl}$  was apparent by the second year of our study and had increased in magnitude by the third year. The physiological and environmental factors leading to  $A_{accl}$  at high  $CO_2$  have been the focus of much experimental and theoretical work (Sage 1994, Lloyd and Farquhar 1996). One factor consistently associated with  $A_{accl}$  in both woody and herbaceous species is a decline in leaf tissue N concentration (McGuire et al. 1995, Peterson et al. 1999), although not all plants with lower leaf N at high  $CO_2$  show strong  $A_{accl}$  (e.g., Will and Ceulemans 1997). It is important in that regard to distinguish between a net reduction in N per unit leaf mass or leaf area that is attributable to a reduction in N content, and a reduction in N concentration due to the dilution effect of leaf starch accumulation without a change in N content. Reduced Rubisco (Ribulose bis-phosphate carboxylase-oxygenase) content is generally the assumed consequence of lower leaf N and is considered a primary physiological mechanism by which  $A_{accl}$  occurs (Stitt 1991). However, plants may compensate for lower leaf-N concentration (mass basis) with additional mesophyll cells, thereby maintaining or even increasing photosynthetic capacity per unit leaf area (Luo et al. 1994).

FIG. 5. Vertical distribution of leaf area in *Populus tremuloides* after 2.5 seasons of growth in soil of low or high N availability and at ambient (open bars) or elevated (shaded bars) CO<sub>2</sub>. The percentage increase due to CO<sub>2</sub> is indicated to the left of the shaded bars (\*  $P < 0.05$ ).



Our results support the model of Luo et al. (1994) identifying reciprocal changes in  $N_m$  (leaf N on a mass basis) and SLA (specific leaf area) as important controls over the direction and magnitude of CO<sub>2</sub> effects on  $A$ . In high-N soil, the modest decline in  $N_m$  is entirely compensated for by increasing leaf thickness and lower SLA, with the result that  $N_a$  (leaf N on an area basis) remains unchanged in elevated- compared to ambient-CO<sub>2</sub>-grown plants, and there is no  $A_{acc}$ . With a greater reduction in  $N_m$  as seen on low-N soil, there was only partial compensation by lower SLA in CO<sub>2</sub>-enriched plants and therefore  $N_a$  declined, leading to significant  $A_{acc}$ . Our data also suggest that the fundamental A:N relationship, reflecting leaf-level allocation of N into different photosynthetic and non-photosynthetic constituents, was unchanged by either CO<sub>2</sub> or soil-N treatments. That is, at a common  $C_i$  (leaf internal CO<sub>2</sub> partial pressure) of 56 Pa, all treatments fell along a common line relating  $\log(A_m)$  ( $A$  on a leaf mass basis) to  $\log(N_m)$ , and whose slope (1.31) was similar to that reported by Reich et al. (1997) for a collection of 111 species across six biomes (1.41). The effect of CO<sub>2</sub> enrichment was to change the intercept of that line, but not its slope. This is equivalent to an increase in photosynthetic nitrogen-use efficiency (Peterson et al. 1999). At equal leaf  $N_m$ , high-CO<sub>2</sub>-grown plants in our study would have a 93% greater  $A_m$ . The acquisition of soil N and its allocation within the plant, both temporally and spatially, is thus clearly of central importance in understanding the long-term consequences of rising CO<sub>2</sub> on photosynthesis in *Populus tremuloides*.

Aboveground biomass accumulation in *P. tremuloides* after 2.5-growing-seasons exposure to elevated CO<sub>2</sub> in soils of contrasting N availability was very similar to the average response of 21 woody-plant species to similar treatments as estimated in the meta-analysis of Curtis and Wang (1998). This was true both in the absolute magnitude of the CO<sub>2</sub> response and in the relative effects of low N availability. We found a 34% increase in stem dry mass at twice-ambient CO<sub>2</sub> and in high-N soil, compared to a 39% increase reported by Curtis and Wang (1998). Low soil-N availability reduced the CO<sub>2</sub> effect on leaf-area growth to

near zero and reduced the stem dry mass responses by ~60%, both common although not universal responses among woody plants (Lloyd and Farquhar 1996). Whole-plant biomass mirrored these responses (+37% at high N, +17% at low N, Zak et al. 2000a) and were again very close to the average values for woody plants estimated by Curtis and Wang (1998). A complete account of CO<sub>2</sub> effects on biomass partitioning in this experiment can be found in the companion paper by Zak et al. (2000a).

Leaf-area distribution within the canopy can have important effects on light interception and growth in *Populus* and can be used in selection programs for rapidly growing genotypes in short-rotation forestry (Ceulemans et al. 1990). The sylleptic growth of lateral branches characteristic of this genus also results in a high degree of architectural plasticity in response to environmental factors such as planting density (Nelson et al. 1981) or soil fertility (Heilman and Xie 1994). In a CO<sub>2</sub>-enrichment study with six tropical tree species of indeterminate growth habit, Reekie and Bazzaz (1989) found that mean canopy height increased at high CO<sub>2</sub> in two species, decreased in one species, and was unchanged in three others. They concluded that CO<sub>2</sub> effects on canopy structure played a major role in competitive interactions among these species. We found little evidence for changes in the vertical distribution of LA (total leaf area per plant) due to growth at high CO<sub>2</sub>. In high-N soil, the majority of LA was in the middle stratum, regardless of CO<sub>2</sub> treatment and consistent with the architecture of *Populus* canopies when grown under similar planting densities (Ceulemans et al. 1990, Kubiske et al. 1997). However, elevated CO<sub>2</sub> caused a greater increase in LA in the upper canopy (+52%) than in the middle (+21%) or lower (+26%) strata, an effect that over time would result in increased self-shading within the canopy and a decline in canopy growth efficiency. Plants grown in low-N soil had much lower total LA relative to plants in high-N soil and the greatest fraction of canopy LA was in the lowest stratum. Again, exposure to high CO<sub>2</sub> did not change this pattern of LA distribution. It should be noted that although our removal of terminal shoot growth during




FIG. 6. Aboveground growth efficiency ( $E$ ; growth efficiency, wood production per unit leaf area per unit time) in relation to chamber leaf-area index (LAI) of *Populus tremuloides* grown in low-N (open symbols) or high-N (solid symbols) soil and at ambient (circles) or elevated (squares)  $\text{CO}_2$ . Growth efficiency was estimated for plants growing in the center of each chamber on the basis of annual stem volume increment ( $E_v$ , symbols without +) or harvested stem mass ( $E_{dm}$ , symbols with +). Data are means  $\pm$  1 SE;  $n = 5$  replicate chambers per  $\text{CO}_2$  and soil-N-availability treatment.

the second winter likely resulted in increased lateral branch production and relatively greater LA lower in the canopy, these effects should be independent of either  $\text{CO}_2$  or soil N treatments since all plants were similarly treated.

*Populus tremuloides* shows substantial genotypic variation in a number of growth and physiological characteristics, including trunk morphology, lateral-branch abscission, timing of leaf drop (Barnes 1959), and sensitivity to ozone stress (Karnosky et al. 1989). Indeed, the large amount of genetic variation present in this, and other, members of the genus *Populus* has provided ample opportunity for breeding and improvements in biomass yield (Farmer 1996). We also observed significant genotypic variation in growth characteristics and in responses to changing soil-N availability. However, we found no genotype  $\times$   $\text{CO}_2$  interactions, indicating a uniform growth response to  $\text{CO}_2$  enrichment by all six genotypes in our study. This is perhaps not surprising given the small number of genotypes examined and the relatively short duration of the experiment. We have found  $\text{CO}_2 \times$  genotype interactions in these same genotypes for condensed tannin production (Mansfield et al. 1999) and early season photosynthetic rates (X. Wang, unpublished data), both of which might eventually influence biomass accumulation. There has been little other work conducted on intraspecific variation in  $\text{CO}_2$  responses within woody plant species. Radoglou and Jarvis (1990) reported variation in leaf structural characteristics at high  $\text{CO}_2$  among hybrid poplar genotypes, while Kalina and Ceulemans (1997) found greater  $A_{acc1}$  at high  $\text{CO}_2$  in a slow-growing poplar genotype compared to a fast-growing genotype. Rising atmospheric  $\text{CO}_2$  clearly has the potential to act as a

selective agent (Curtis et al. 1996) and hence alter both ecosystem composition and function. Identification of traits that are responsive to  $\text{CO}_2$  enrichment and that contribute directly to growth and fitness should receive continued attention by forest geneticists and evolutionary biologists.

Whole-plant net C gain is a function of the assimilatory capacity of leaves, the area and arrangement in space and time of leaf surface, and the loss of carbon through respiration, tissue abscission, and processes such as rhizodeposition and volatile organic-carbon emissions. Elevated  $\text{CO}_2$  has the potential to affect each of these processes, and it is therefore not surprising that there is no simple relationship between, for example,  $\text{CO}_2$  effects on  $A$  and  $\text{CO}_2$  effects on harvested biomass. Measures that integrate assimilatory and allocational processes over time, such as those obtained through growth analysis, may be more useful in interpreting and predicting environmental effects on production than are short-term physiological measures such as  $A$ . In forests, as with herbaceous crops, productivity is strongly related to total light interception, and stand growth, or yield, may be estimated by a knowledge of maximum canopy leaf-area index (LAI) and growth efficiency ( $E$ , wood production per unit leaf area per unit time) (Waring 1983). As stand development proceeds,  $E$  typically declines due to increased competition for light and allocation to non-photosynthetic tissue (Cannell 1989). However, productivity will continue to increase as long as LAI increases faster than  $E$  decreases. Heilman and Xie (1994) found a linear relationship between stand productivity and LAI in poplar hybrids up to an LAI of  $\sim 6$ , with a declining rate of increase at higher LAI.

Growth efficiency was low in our experiment during the first year of growth, likely due to the energy costs of root establishment following outplanting from cuttings. Although we could not measure total belowground biomass prior to harvest, a decline in root-to-shoot ratio with plant age, as is typical in young *Populus* plants (Sheppard and Smith 1993), would result in increasing  $E$  with time, as was observed in our study. This trend would be opposed by increasing competition for light as LAI increased and by greater sapwood respiratory demand with increasing stem volume. Neither soil-N availability nor  $\text{CO}_2$  enrichment had any apparent effect on the relationship between LAI and  $E_v$ , however. That is, while at any point in time plants in high-N soil were larger than plants in low-N soil, and  $\text{CO}_2$  enrichment increased plant size in high-N soil, at equivalent stages of canopy development all treatments produced stem volume with equal efficiency. The basic allometric relationship between LA and stem volume (or mass) increment was unaltered by  $\text{CO}_2$  enrichment, and we would therefore expect no change in aboveground stand wood volume or mass at equivalent LAIs, at least through an LAI of  $\sim 6$ . These results support

others showing little effect of elevated CO<sub>2</sub> on woody-plant allometry (Gebauer et al. 1996, Zak et al. 2000a).

However, important questions remain. In particular, we do not know whether  $E$  will remain equivalent between CO<sub>2</sub> treatments as LAI increases. If quantum yield is increased by high CO<sub>2</sub>, as suggested by Long and Drake (1991), we would expect  $E$  to be sustained as LAI increases to a greater extent in elevated- compared to ambient-CO<sub>2</sub>-grown plants due to reduced self-shading effects on carbon assimilation. We also do not know maximum LAI for any of the treatment combinations, another important determinant of long-term stand productivity. For example, maximum LAI in low-N soil or at ambient CO<sub>2</sub> may never reach that obtained in high-N soil or at elevated CO<sub>2</sub>. Answers to these questions require experiments at temporal and spatial scales beyond those achievable with open-top chambers.

These results illustrate several important issues regarding tree growth at high CO<sub>2</sub>. The first is that increased  $A_{gr}$  does not necessarily lead to increased  $E$ . For *P. tremuloides* in high-N soil, higher  $A_{gr}$  due to CO<sub>2</sub> enrichment was offset by higher rates of carbon loss, via increased leaf and stem respiration and from greater fine-root turnover (Pregitzer et al. 2000). In effect, increased stem-biomass accumulation could be accounted for entirely by greater LA. In low-N soil, CO<sub>2</sub> stimulation of  $A_{gr}$  was less, and varied between years, but also did not translate into significantly greater  $E$ . The second issue is that knowledge of the dynamic response of  $E$  with respect to LAI will be necessary to answer the question of whether trees will grow larger or simply grow faster during stand development (Eamus and Jarvis 1989). Our data suggest the latter. That is, in high-N soil stand productivity increased at high CO<sub>2</sub>, but the fundamental relationship between tree growth and canopy development appeared unchanged (see also Zak et al. 2000a).

#### Summary and implications

Elevated atmospheric CO<sub>2</sub> had a sustained, positive effect on carbon-gain capacity and aboveground biomass accumulation in *P. tremuloides*, but only under conditions of high soil-N availability. Carbon dioxide enrichment increased photosynthetic N-use efficiency regardless of soil-N availability but in low-N soil growth at elevated CO<sub>2</sub> resulted in lower leaf-N concentration. This led to a reduction in photosynthetic capacity relative to plants at ambient CO<sub>2</sub>. Gains in net CO<sub>2</sub>-assimilation rate under elevated CO<sub>2</sub> were largely offset by increased dark respiration and fine-root turnover, resulting in no net increase in growth efficiency at high CO<sub>2</sub>. Nonetheless, increased stand productivity with elevated CO<sub>2</sub> and high soil-N availability could shorten harvest rotation intervals for this economically important species and speed ecological succession in aggrading northern hardwood forests.

#### ACKNOWLEDGMENTS

We thank Diana Randlett and Bill Holmes for help in experimental design and execution, Mark Kalnins for leaf anatomical work, and Richard Norby for comments on an earlier draft. This work was funded through grants from the National Institute for Global Environmental Change (DOE-NIGEC), Program for Ecosystem Research (DOE-PER Grant DE-FGOZ-93ER6166), the University of Michigan Project for the Integrated Study of Global Change, and the University of Michigan Biological Station.

#### LITERATURE CITED

- Barnes, B. V. 1959. Natural variation and clonal development of *Populus tremuloides* and *P. grandidentata* in northern lower Michigan. Dissertation. University of Michigan, Ann Arbor, Michigan, USA.
- Berntson, G., and F. Bazzaz. 1997. Nitrogen cycling in microcosms of yellow birch exposed to elevated CO<sub>2</sub>: simultaneous positive and negative below-ground feedbacks. *Global Change Biology* 3:247–258.
- Cannell, M. G. R. 1989. Light interception, light use efficiency and assimilate partitioning in poplar and willow stands. Pages 1–12 in J. S. Pereira and J. J. Landsberg, editors. Biomass production by fast-growing trees. Kluwer Academic, Dordrecht, The Netherlands.
- Ceulemans, R., X. N. Jiang, and B. Y. Shao. 1995. Effects of elevated CO<sub>2</sub> on growth, biomass production and nitrogen allocation of two *Populus* clones. *Journal of Biogeography* 22:261–268.
- Ceulemans, R., R. F. Stettler, T. M. Hinckley, J. G. Isebrands, and P. E. Heilman. 1990. Crown architecture of *Populus* clones as determined by branch orientation and branch characteristics. *Tree Physiology* 7:157–167.
- Cooper, H., and L. V. Hedges. 1994. The handbook of research synthesis. Russell Sage Foundation, New York, New York, USA.
- Curtis, P. S. 1996. A meta-analysis of leaf gas exchange and nitrogen in trees grown under elevated carbon dioxide. *Plant Cell and Environment* 19:127–137.
- Curtis, P. S., D. Jenkins Klus, S. Kalisz, and S. J. Tonsor. 1996. Intra-specific variation in CO<sub>2</sub> responses in *Raphanus raphanistrum* and *Plantago lanceolata*: assessing the potential for evolutionary change with rising atmospheric CO<sub>2</sub>. Pages 13–22 in Ch. Körner and F. A. Bazzaz, editors. Carbon dioxide, populations, and communities. Academic Press, San Diego, California, USA.
- Curtis, P. S., E. G. O'Neill, J. A. Teeri, D. R. Zak, and K. S. Pregitzer. 1994. Belowground responses to rising atmospheric CO<sub>2</sub>: implications for plants, soil biota, and ecosystem processes. *Plant and Soil* 165:1–6.
- Curtis, P. S., C. S. Vogel, K. S. Pregitzer, D. R. Zak, and J. A. Teeri. 1995. Interacting effects of soil fertility and atmospheric CO<sub>2</sub> on leaf area growth and carbon gain physiology in *Populus × euramericana* (Dode) Guinier. *New Phytologist* 129:253–263.
- Curtis, P. S., and X. Wang. 1998. A meta-analysis of elevated CO<sub>2</sub> effects on woody plant mass, form, and physiology. *Oecologia* 113:299–313.
- Diaz, S., J. P. Grime, J. Harris, and E. McPherson. 1993. Evidence of a feedback mechanism limiting plant response to elevated carbon dioxide. *Nature* 364:616–617.
- Eamus, D., G. A. Duff, and C. A. Berryman. 1995. Photosynthetic responses to temperature, light flux density, CO<sub>2</sub> concentration and vapour pressure deficit in *Eucalyptus tetrodonta* grown under CO<sub>2</sub> enrichment. *Environmental Pollution* 90:41–49.
- Eamus, D., and P. G. Jarvis. 1989. Direct effects of CO<sub>2</sub> increases on trees and forests (natural and commercial) in the UK. *Advances in Ecological Research* 19:1–55.
- Farmer, R. E. J. 1996. The genealogy of *Populus*. Pages

- 33–56 in R. F. Stettler, H. D. J. Bradshaw, P. E. Heilman, and T. M. Hinckley, editors. *Biology of Populus and its implications for management and conservation*. NRC Research Press, Ottawa, Ontario, Canada.
- Field, C., and H. A. Mooney 1986. The photosynthesis–nitrogen relationship in wild plants. Pages 25–55 in T. J. Givnish, editor. *On the economy of plant form and function*. Cambridge University Press, New York, New York, USA.
- Gebauer, R. L. E., J. E. Reynolds, and B. R. Strain. 1996. Allometric relations and growth in *Pinus taeda*: the effect of elevated CO<sub>2</sub> and changing N availability. *New Phytologist* **134**:85–93.
- Gunderson, C. A., R. J. Norby, and S. D. Wullschlegel. 1993. Foliar gas exchange responses of two deciduous hardwoods during 3 years of growth in elevated CO<sub>2</sub>: no loss of photosynthetic enhancement. *Plant Cell and Environment* **16**:797–807.
- Gunderson, C. A., and S. D. Wullschlegel. 1994. Photosynthetic acclimation in trees to rising atmospheric CO<sub>2</sub>: a broader perspective. *Photosynthesis Research* **39**:369–388.
- Heagle, A. S., R. B. Philbeck, R. E. Ferrell, and W. W. Heck. 1989. Design and performance of a large, field exposure chamber to measure effects of air quality on plants. *Journal of Environmental Quality* **18**:361–368.
- Hedges, L. V., J. Gurevitch, and P. S. Curtis. 1999. The meta-analysis of response ratios in experimental ecology. *Ecology* **80**:1150–1156.
- Heilman, P. E., and F. Xie. 1994. Effects of nitrogen fertilizer on leaf area, light interception, and productivity of short-rotation *Populus trichocarpa* × *Populus deltoides* hybrids. *Canadian Journal of Forest Research* **24**:166–173.
- Hungate, B. A., E. A. Holland, R. B. Jackson, F. S. Chapin III, C. B. Field, and H. A. Mooney. 1997. The fate of carbon in grasslands under carbon dioxide enrichment. *Nature* **388**:576–579.
- Idso, K. E., and S. B. Idso. 1994. Plant responses to atmospheric CO<sub>2</sub> enrichment in the face of environmental constraints: a review of the past 10 years' research. *Agricultural and Forest Meteorology* **69**:153–203.
- Jones, M. G. K., W. H. J. Outlaw, and O. H. Lowry. 1977. Enzymic assay of 10(–7) to 10(–14) moles of sucrose in plant tissues. *Plant Physiology* **60**:379–383.
- Kalina, J., and R. Ceulemans. 1997. Clonal differences in the response of dark and light reactions of photosynthesis to elevated atmospheric CO<sub>2</sub> in poplar. *Photosynthetica* **33**:51–61.
- Karnosky, D. F., P. C. Berrang, F. Scholz, and J. P. Bennett. 1989. Variation in and natural selection for air pollution tolerances in trees. Pages 29–37 in F. Scholz, H.-R. Gregorius, and D. Rudin, editors. *Genetic effects of air pollutants in forest tree populations*. Springer-Verlag, Berlin, Germany.
- Kubiske, M. E., K. S. Pregitzer, C. J. Mikan, D. R. Zak, J. L. Maziasz, and J. A. Teeri. 1997. *Populus tremuloides* photosynthesis and crown architecture in response to elevated CO<sub>2</sub> and soil N availability. *Oecologia* **110**:328–336.
- Kuo, S. 1996. Phosphorus. Pages 894–895 in D. L. Sparks, A. L. Page, P. A. Helmke, R. H. Loeppert, P. N. Soltanpour, M. A. Tabatabai, C. T. Johnston, and M. E. Summer, editors. *Methods of soil analysis*. Part 3. Chemical methods. SSSA Book Series number 5. Soil Science Society of America and American Society of Agronomy, Madison, Wisconsin, USA.
- Lambers, H., and H. Poorter. 1992. Inherent variation in growth rate between higher plants: a search for physiological causes and ecological consequences. *Advances in Ecological Research* **23**:187–261.
- Lewis, J. D., D. T. Tissue, and B. R. Strain. 1996. Seasonal response of photosynthesis to elevated CO<sub>2</sub> in loblolly pine (*Pinus taeda* L.) over two growing seasons. *Global Change Biology* **2**:103–114.
- Lloyd, J., and G. D. Farquhar. 1996. The CO<sub>2</sub> dependence of photosynthesis, plant growth responses to elevated atmospheric CO<sub>2</sub> concentrations and their interaction with soil nutrient status. I. General principles and forest ecosystems. *Functional Ecology* **10**:4–32.
- Long, S. P., and B. G. Drake. 1991. Effect of the long-term elevation of CO<sub>2</sub> concentration in the field on the quantum yield of photosynthesis of the C<sub>3</sub> sedge, *Scirpus olney* L. *Plant Physiology* **96**:221–226.
- Luo, Y., C. B. Field, and H. A. Mooney. 1994. Predicting responses of photosynthesis and root fraction to elevated [CO<sub>2</sub>]: interactions among carbon, nitrogen, and growth. *Plant Cell and Environment* **17**:1195–1204.
- Mansfield, J. W., P. S. Curtis, D. R. Zak, and K. S. Pregitzer. 1999. Genotypic variation for condensed tannin production in trembling aspen (*Populus tremuloides*) under elevated CO<sub>2</sub> and in high- and low-fertility soil. *American Journal of Botany* **86**:1154–1159.
- McConnaughay, K. D. M., G. M. Berntson, and F. A. Bazzaz. 1993. Plant responses to carbon dioxide. *Nature* **361**:24.
- McGill, W. B., K. R. Cannon, J. A. Robertson, and F. D. Cook. 1986. Dynamics of soil microbial biomass and water-soluble organic C in Breton L after 50 years of cropping to 2 rotations. *Canadian Journal of Soil Science* **66**:1–19.
- McGuire, A. D., J. M. Melilo, and L. A. Joyce. 1995. The role of nitrogen in the response of forest net primary production to elevated atmospheric carbon dioxide. *Annual Review of Ecology and Systematics* **26**:473–504.
- Nelson, N. D., T. Burk, and J. G. Isebrands. 1981. Crown architecture of short-rotation, intensively cultured *Populus*. I. Effects of clone and spacing on first-order branch characteristics. *Canadian Journal of Forest Research* **11**:73–81.
- Norby, R. J. 1996. Forest canopy productivity index. *Nature* **381**:564.
- Norby, R. J., C. A. Gunderson, S. D. Wullschlegel, E. G. O'Neill, and M. K. McCracken. 1992. Productivity and compensatory responses of yellow-poplar trees in elevated CO<sub>2</sub>. *Nature* **357**:322–324.
- Norby, R. J., S. D. Wullschlegel, C. A. Gunderson, D. W. Johnson, and R. Ceulemans. 1999. Tree responses to rising CO<sub>2</sub> in field experiments: implications for the future forest. *Plant Cell and Environment* **22**:683–714.
- Pastor, J., and W. M. Post. 1988. Response of northern forests to CO<sub>2</sub>-induced climate change. *Nature* **334**:55–58.
- Peterson, A. G., J. T. Ball, Y. Luo, C. B. Field, P. B. Reich, P. S. Curtis, K. L. Griffin, C. A. Gunderson, R. J. Norby, D. T. Tissue, M. Forstreuter, A. Rey, C. S. Vogel, and CMEAL participants. 1999. The photosynthesis–leaf nitrogen relationship at ambient and elevated atmospheric carbon dioxide: a meta-analysis. *Global Change Biology* **5**:331–346.
- Pregitzer, K. S., D. R. Zak, J. Maziasz, J. DeForest, P. S. Curtis, and J. Lussenhop. 2000. Interactive effects of atmospheric CO<sub>2</sub> and soil-N availability on the fine roots of *Populus tremuloides*. *Ecological Applications* **10**:18–33.
- Radoglou, K. M., and P. G. Jarvis. 1990. Effects of CO<sub>2</sub> enrichment on four poplar clones. I. Growth and leaf anatomy. *Annals of Botany* **65**:617–626.
- Reekie, E. G., and F. A. Bazzaz. 1989. Competition and patterns of resource use among seedlings of five tropical trees grown at ambient and elevated CO<sub>2</sub>. *Oecologia* **79**:212–222.
- Reich, P. B., M. B. Walters, and D. S. Ellsworth. 1997. From tropics to tundra: global convergence in plant functioning. *Proceedings of the National Academy of Sciences (USA)* **94**:13730–13734.
- Reynolds, J. F., P. R. Kemp, B. Acock, J.-L. Chen, and D. L. Moorhead. 1996. Progress, limitations, and challenges in

- modeling the effects of elevated CO<sub>2</sub> on plants and ecosystems. Pages 347–380 in G. W. Koch and H. A. Mooney, editors. Carbon dioxide and terrestrial ecosystems. Academic Press, New York, New York, USA.
- Sage, R. F. 1994. Acclimation of photosynthesis to increasing atmospheric CO<sub>2</sub>: the gas-exchange perspective. *Photosynthesis Research* **39**:351–368.
- Sheppard, W. D., and F. W. Smith. 1993. The role of near-surface lateral roots in the life cycle of aspen in the central Rocky Mountains. *Forest Ecology and Management* **61**: 157–170.
- Sokal, R. R., and F. J. Rohlf 1981. *Biometry*. W. H. Freeman & Company, San Francisco, California, USA.
- Stitt, M. 1991. Rising CO<sub>2</sub> levels and their potential significance for carbon flow in photosynthetic cells. *Plant Cell and Environment* **14**:741–762.
- Thomas, R. B., and B. R. Strain. 1991. Root restriction as a factor in photosynthetic acclimation of cotton seedlings grown in elevated carbon dioxide. *Plant Physiology* **96**: 629–634.
- Tissue, D. T., R. B. Thomas, and B. R. Strain. 1993. Long-term effects of elevated CO<sub>2</sub> and nutrients on photosynthesis and rubisco in loblolly pine seedlings. *Plant Cell and Environment* **16**:859–865.
- Vitousek, P. M., and P. A. Matson. 1984. Mechanisms of nitrogen retention in forest ecosystems: a field experiment. *Science* **225**:51–52.
- Waring, R. H. 1983. Estimating forest growth and efficiency in relation to canopy leaf area. *Advances in Ecological Research* **13**:327–354.
- Will, R. E., and R. Ceulemans. 1997. Effects of elevated CO<sub>2</sub> concentration on photosynthesis, respiration and carbohydrate status on coppice *Populus* hybrids. *Physiologia Plantarum* **100**:933–939.
- Wullschlegel, S. D., W. M. Post, and A. W. King. 1995. On the potential for a CO<sub>2</sub> fertilization effect in forests: estimates of the biotic growth factor based on 58 controlled-exposure studies. Pages 85–107 in A. W. Woodwell and F. T. Mackensie, editors. Biospheric feedbacks in the global climate system: Will warming feed the warming? Oxford University Press, London, UK.
- Zak, D. R., D. F. Grigal, S. Gleeson, and D. Tilman. 1990. Carbon and nitrogen cycling during secondary succession: constraints on plant and microbial productivity. *Biogeochemistry* **11**:111–129.
- Zak, D. R., K. S. Pregitzer, P. S. Curtis, and W. E. Holmes. 2000a. Atmospheric CO<sub>2</sub> and the composition and function of soil microbial communities. *Ecological Applications* **10**: 47–59.
- Zak, D. R., K. S. Pregitzer, P. S. Curtis, J. A. Teeri, R. Fogel, and D. L. Randlett. 1993. Elevated atmospheric CO<sub>2</sub> and feedback between carbon and nitrogen cycles. *Plant and Soil* **151**:105–117.
- Zak, D. R., K. S. Pregitzer, P. S. Curtis, C. S. Vogel, W. E. Holmes, and J. Lussenhop. 2000b. Atmospheric CO<sub>2</sub>, soil-N availability, and allocation of biomass and nitrogen by *Populus tremuloides*. *Ecological Applications* **10**:34–46.