

# Tropospheric O<sub>3</sub> compromises net primary production in young stands of trembling aspen, paper birch and sugar maple in response to elevated atmospheric CO<sub>2</sub>

John S. King<sup>1,\*</sup>, Mark E. Kubiske<sup>2</sup>, Kurt S. Pregitzer<sup>1</sup>, George R. Hendrey<sup>3</sup>, Evan P. McDonald<sup>4</sup>, Christian P. Giardina<sup>5</sup>, Vanessa S. Quinn<sup>2</sup> and David F. Karnosky<sup>1</sup>

<sup>1</sup>Ecosystem Science Center, School of Forest Resources and Environmental Science, Michigan Technological University, Houghton, Michigan 49931, USA;

<sup>2</sup>USDA Forest Service, North-central Research Station, Rhinelander, Wisconsin 54501, USA; <sup>3</sup>Brookhaven National Laboratory, Department of Environmental

Science, Earth System Sciences Division, Upton, New York 11973, USA; <sup>4</sup>Department of Forest Ecology and Management, University of Wisconsin-Madison,

Madison, Wisconsin 53706, USA; <sup>5</sup>USDA Forest Service, North-central Research Station, Houghton, Michigan 49931, USA; \*Present address: Department

of Forestry and Environmental Resources, North Carolina State University, Raleigh, NC 27695, USA

## Summary

Author for correspondence:

John S. King

Tel: +1 919 513 7855

Fax: +1 919 515 3169

Email: john\_king@ncsu.edu

Received: 20 May 2005

Accepted: 8 July 2005

- Concentrations of atmospheric CO<sub>2</sub> and tropospheric ozone (O<sub>3</sub>) are rising concurrently in the atmosphere, with potentially antagonistic effects on forest net primary production (NPP) and implications for terrestrial carbon sequestration.
- Using free-air CO<sub>2</sub> enrichment (FACE) technology, we exposed north-temperate forest communities to concentrations of CO<sub>2</sub> and O<sub>3</sub> predicted for the year 2050 for the first 7 yr of stand development. Site-specific allometric equations were applied to annual nondestructive growth measurements to estimate above- and below-ground biomass and NPP for each year of the experiment.
- Relative to the control, elevated CO<sub>2</sub> increased total biomass 25, 45 and 60% in the aspen, aspen–birch and aspen–maple communities, respectively. Tropospheric O<sub>3</sub> caused 23, 13 and 14% reductions in total biomass relative to the control in the respective communities. Combined fumigation resulted in total biomass response of –7.8, +8.4 and +24.3% relative to the control in the aspen, aspen–birch and aspen–sugar maple communities, respectively.
- These results indicate that exposure to even moderate levels of O<sub>3</sub> significantly reduce the capacity of NPP to respond to elevated CO<sub>2</sub> in some forests.

**Key words:** Aspen FACE (free-air CO<sub>2</sub> enrichment), elevated carbon dioxide, global change, net primary production (NPP), tropospheric ozone (O<sub>3</sub>).

*New Phytologist* (2005) **168**: 623–636

© The Authors (2005). Journal compilation © *New Phytologist* (2005)

doi: 10.1111/j.1469-8137.2005.01557.x

## Introduction

Land-based and remotely sensed data show a carbon sink in Northern Hemisphere forests of 0.30–0.68 petagrams (Pg = 10<sup>15</sup> g) per year for the 1980s and 1990s, with 70% occurring in Eurasia (Myneni *et al.*, 2001; Pacala *et al.*, 2001). In the USA, the forest carbon sink was expanding because of forest recovery on former agricultural land and fire suppression, which offset 10–30% of USA fossil fuel emissions during the 1980s (Houghton *et al.*, 1999; Caspersen *et al.*,

2000; Birdsey & Lewis, 2002). Rising atmospheric CO<sub>2</sub> has received considerable attention as a possible stimulus to forest net primary production (NPP) that could further offset fossil fuel emissions (Ceulemans & Mousseau, 1994; Curtis & Wang, 1998; Houghton *et al.*, 2001). Less well recognized, however, is that tropospheric O<sub>3</sub> is also increasing globally (Fowler *et al.*, 1999), and is probably reducing the potential enhancement of forest NPP and terrestrial C sequestration caused by elevated atmospheric CO<sub>2</sub> (Karnosky *et al.*, 1999, 2003; Loya *et al.*, 2003; Felzer *et al.*, 2004).

Decades of experimental evidence show that small forest trees experience an average stimulation of 16–31% in biomass production under elevated atmospheric CO<sub>2</sub>, but these responses can be constrained by soil nutrient availability or other environmental factors (Strain & Cure, 1994; McGuire *et al.*, 1995; Gebauer *et al.*, 1996; Curtis & Wang, 1998; Johnson, 1999; Zak *et al.*, 2000; Oren *et al.*, 2001). Evidence from long-term forest FACE (free-air CO<sub>2</sub> enrichment) experiments corroborates this finding at larger spatial and temporal scales. Net primary production of a 13-yr-old loblolly pine (*Pinus taeda*) ecosystem was stimulated 26% under CO<sub>2</sub> enrichment, and this response persisted for 4 yr (DeLucia *et al.*, 1999; Hamilton *et al.*, 2002). A 10-yr-old sweetgum (*Liquidambar styraciflua*) ecosystem increased NPP an average 21% under CO<sub>2</sub> enrichment (Norby *et al.*, 2002). Three species of *Populus* grown in short rotation culture increased woody biomass by 15–27% under elevated CO<sub>2</sub> (Calfapietra *et al.*, 2003). Responses of mature forests to elevated atmospheric CO<sub>2</sub> are still poorly understood.

Most elevated CO<sub>2</sub> studies have not considered the phytotoxic effects of simultaneous exposure to elevated tropospheric O<sub>3</sub>, which has been shown to decrease tree seedling growth from –2 to –69% (Pye, 1988). Pre-industrial concentrations of tropospheric O<sub>3</sub> are estimated to have been less than 10 nl l<sup>-1</sup>, and have risen to 30–40 nl l<sup>-1</sup> background levels today (Levy *et al.*, 1997). Tropospheric O<sub>3</sub> concentrations are expected to exceed 60 nl l<sup>-1</sup> over large portions (50%) of the global forested land surface by the year 2100 (Felzer *et al.*, 2004). The decrease in agricultural productivity caused by ambient O<sub>3</sub> toxicity has been estimated at US\$1.0–5.8 billion annually (1990 dollars) in the USA (Kopp *et al.*, 1985; Adams *et al.*, 1986; Murphy *et al.*, 1999), but economic losses to wood production are unknown (Krupa *et al.*, 2000). Biogeochemical modeling estimates of reductions in annual terrestrial C sequestration in the USA, caused by ambient O<sub>3</sub> pollution during the late 1980s to early 1990s, range from 18 to 38 Tg C yr<sup>-1</sup>, which must be accounted for in future calculations of the global C budget (Felzer *et al.*, 2004). Therefore a key to understanding the role of forests in

mitigating the build-up of atmospheric CO<sub>2</sub> is determining how NPP will respond to the interactive effects of elevated atmospheric CO<sub>2</sub> and tropospheric O<sub>3</sub>.

Here we report on the NPP of intact experimental forest communities dominated by the most widespread tree taxa in North America, in response to the interactive effects of elevated atmospheric CO<sub>2</sub> and tropospheric O<sub>3</sub>. The study was performed at the Aspen FACE project in Rhineland, WI, USA (Dickson *et al.*, 2000). Communities of pure trembling aspen (*Populus tremuloides* Michx.), and competitive mixes of trembling aspen–paper birch (*Betula papyrifera* Marsh.) and trembling aspen–sugar maple (*Acer saccharum* Marsh.) were exposed for 7 yr to concentrations of atmospheric CO<sub>2</sub> and tropospheric O<sub>3</sub> predicted for the year 2050. Net primary production was estimated with species-specific allometric regressions developed at the site, applied to annual nondestructive measurements of all trees in the experimental plots. We hypothesized that (i) elevated atmospheric CO<sub>2</sub> would provide sustained enhancement of NPP in all three forest communities; and (ii) elevated tropospheric O<sub>3</sub> would decrease it. Because it has been postulated that elevated CO<sub>2</sub> may decrease O<sub>3</sub> uptake into the plant because of decreased stomatal conductance (Allen, 1990; Wustman *et al.*, 2003), our third hypothesis was that simultaneous exposure to both elevated CO<sub>2</sub> and elevated tropospheric O<sub>3</sub> would result in growth similar to the control.

## Materials and Methods

### Field experiment

The Aspen FACE project is a randomized complete block design of atmospheric CO<sub>2</sub> and tropospheric O<sub>3</sub> treatments, with species composition (aspen, aspen–birch, and aspen–maple) as a split-plot factor ( $n = 3$ ). The 12 30-m-diameter plots are fumigated using free-air technology of the Brookhaven National Laboratory's design (Hendrey *et al.*, 1999) to maintain atmospheric targets of 560 ppm CO<sub>2</sub> and 1.5× ambient O<sub>3</sub> (Table 1). Fumigation is during daylight hours only, and begins at bud break in the spring and ends at

Treatment	1998	1999	2000	2001	2002	2003
Ambient CO <sub>2</sub> (control, +O <sub>3</sub> ) (ppm)	343	347	347	356	361	367
Elevated CO <sub>2</sub> (+CO <sub>2</sub> , +CO <sub>2</sub> + O <sub>3</sub> ) (ppm)	530	548	547	528	537	535
Ambient O <sub>3</sub> (control, +CO <sub>2</sub> ) (ppb)	37.5	36.9	36.0	38.8	33.1	38.0
Elevated O <sub>3</sub> (+O <sub>3</sub> , +CO <sub>2</sub> + O <sub>3</sub> ) (ppb)	54.5	51.9	49.3	52.6	49.5	51.0
Ambient O <sub>3</sub> exposure (control, +CO <sub>2</sub> ) (ppmh)	63.8	62.8	58.2	66.1	54.3	60.4
Elevated O <sub>3</sub> exposure (+O <sub>3</sub> , +CO <sub>2</sub> + O <sub>3</sub> ) (ppmh)	97.4	88.8	81.6	90.0	81.4	81.1
Exposure duration (d)	166	143	139	143	138	145

**Table 1** Summary of Aspen FACE project control data for atmospheric fumigation treatments, 1998–2003. Values are mean daily concentrations or sums (O<sub>3</sub> exposure) for the entire growing season of each year

leaf senescence in the fall, with an average growing season of 145 d (Table 1). Trees were planted at 1 × 1-m spacing, and fumigation began in 1997. Over the life of the experiment, system performance has been within 20% of the target 93% of the time for CO<sub>2</sub>, and within 20% of the target 80% of the time for O<sub>3</sub> ([www.aspenface.mtu.edu](http://www.aspenface.mtu.edu)).

Tree growth is monitored by annual measurement of total height and diameter at 10 cm above-ground of all trees within the central core area of the plots ( $n = 3684$ ). The core area in each plot measures 166, 76 and 65 m<sup>2</sup> for the aspen, aspen–birch and aspen–maple sections, respectively. The core area in which nondestructive studies are carried out is buffered from possible edge effects by five rows of trees on the outer edge of the plots. In 2000 and 2002, complete above- and below-ground harvests were performed on a total of 196 trees at the edge of the central core area to develop species-specific allometric biomass equations. Trees were harvested in midsummer at peak leaf area by severing the stem 3 cm above ground, separating foliage from wood (stem + branches). The tree heart root system was sampled by driving a 25.4-cm internal diameter corer to a depth of 25 cm centered on the severed tree stem. Between-tree coarse and fine (<1 mm diameter) root biomass was estimated with 10 randomly located cores, 15 cm diameter by 25 cm depth, within the subplots. All roots were removed from the soil by washing over a fine-mesh screen, and were sorted into coarse (heart + all roots >1 mm diameter) and fine (all roots <1 mm diameter). Only live root data are presented in the current study. All plant parts were dried to constant mass at 65°C and subsamples were combusted at 500°C for 7 h to correct for mineral content.

### Estimating stand-level biomass and NPP

Allometric biomass equations (Table 2) were developed by regressing the logarithm of plant dry weight (g; foliage, wood, heart root system) against the logarithm of stem diameter (cm) of the harvested trees for each species (aspen, paper birch,

sugar maple). Regressions were tested for significant effects of the experimental treatments on model intercept and slope, as well as for suitability of transformations and independent variables. Models of the form  $\ln(\text{dependent variable}) = B_0 + B_1 \ln(\text{diameter})$  proved to explain most of the variation in the data, with statistically insignificant additional predictive power added by height. Allometric analyses of the biomass data from the two harvests (King *et al.*, 1996, 1999) indicated that biomass partitioning between foliage, wood and heart roots was unaffected by the treatments, allowing for use of a common model for each species. Model  $R^2$  values ranged from 0.84 to 0.99, with seven of the nine models having  $R^2 > 0.90$  (Table 2).

The species-specific regressions were applied to the annual diameter measurements of all trees in the core area of the plots for each year of the experiment, correcting for slight underestimation during back-transformation of biomass estimates using the method of Baskerville (1972). Biomass estimates of foliage, wood and heart roots of all trees in the subplots (g) were summed and divided by subplot area to arrive at stand-level estimates of tree biomass (g m<sup>-2</sup>). Between-tree coarse and fine root biomass (g m<sup>-2</sup>) for each year of the experiment was estimated by assuming that the partitioning of total root biomass to heart, coarse and fine root fractions determined by destructive harvest in 2000 and 2002 did not change over time. We then applied the respective root fractions to the allometrically determined estimates of heart root biomass for each year, to arrive at total root biomass. Annual fine root production cannot be determined by this method; however, a published record of annual peak fine root biomass will be useful when fine root turnover rates become available. Additionally, presentation of fine root biomass data allows comparison of the relative size of the fine root pool relative to other plant parts. Annual net wood and coarse root production was determined by subtracting previous year biomass from subsequent year biomass (g m<sup>-2</sup> yr<sup>-1</sup>). Tree mortality was not quantified explicitly in the current study; however, individual trees contributed to stand-level biomass and production estimates

**Table 2** Allometric regressions used to predict tree component biomass of young aspen, paper birch and sugar maple at the Aspen FACE project in Rhinelander, WI, USA

Dependent variable	Intercept (P)	Parameter estimate (P)	MSE	R <sup>2</sup>	n
Aspen foliage	1.48984 (<0.0001)	2.70111 (<0.0001)	0.14649	0.892	131
Aspen wood	1.48067 (<0.0001)	1.87997 (<0.0001)	0.04385	0.929	128
Aspen heart root	2.86029 (<0.0001)	1.87143 (<0.0001)	0.04421	0.929	128
Paper birch foliage	1.78036 (<0.0001)	2.38384 (<0.0001)	0.22940	0.836	37
Paper birch wood	3.19439 (<0.0001)	2.50650 (<0.0001)	0.06087	0.955	37
Paper birch heart root	2.48509 (<0.0001)	1.98989 (<0.0001)	0.05909	0.932	37
Sugar maple foliage	2.35586 (<0.0001)	2.29003 (<0.0001)	0.13239	0.906	25
Sugar maple wood	2.93748 (<0.0001)	2.88168 (<0.0001)	0.06722	0.968	25
Sugar maple heart root	3.03418 (<0.0001)	1.79167 (<0.0001)	0.10574	0.883	24

MSE, mean square error.

Models were developed from trees harvested destructively within FACE plots in 2000 and 2002. All models had the form  $\log(y) = m \log(x) + b$ , where  $y$  = biomass component (g) and  $x$  = diameter (cm). Baskerville's (1972) adjustment to the antilogarithm was applied when calculating absolute data from the log–log models.

only as long as they were alive. At the point when trees were recorded as 'dead' they were removed from subsequent allometric modeling, thereby implicitly accounting for mortality.

Stand-level estimates of total and component biomass and production were tested for main effects ( $\text{CO}_2$ ,  $\text{O}_3$ ) and split-plot effects (community, time) using split-plot ANOVA appropriate for the Aspen FACE experimental design (King *et al.*, 2001) using the STATISTICAL ANALYSIS SYSTEM software (SAS Institute, Cary, North Carolina, USA). Tree size in 1997 (diameter<sup>2</sup> × height) was used as a covariate in the ANOVA to account for initial differences in plant size.


## Results

### Annual biomass production

In 1997, the newly planted stands averaged 22, 15 and 11 g biomass m<sup>-2</sup> in the aspen, aspen–birch and aspen–maple

communities, respectively (Table 3; Fig. 1). Annual production of foliage, wood (stem + branches), and coarse roots in control plots increased from 1998 to 2003 in all communities (time  $P < 0.000$ ). Over this time, annual foliage production averaged 250, 213 and 127 g m<sup>-2</sup> in the aspen, aspen–birch and aspen–maple communities, respectively. Wood production averaged 372, 285 and 190 g m<sup>-2</sup>, while coarse root production averaged 90, 76 and 52 g m<sup>-2</sup> in the aspen, aspen–birch and aspen–maple communities, respectively. Annual fine root production was not quantified in the current study; however, July fine root biomass averaged 24, 20 and 24 g m<sup>-2</sup> in the aspen, aspen–birch and aspen–maple control plots, respectively, from 1998 to 2003.

Annual biomass production of foliage, wood and coarse roots varied in response to the treatments over time, with significant  $\text{CO}_2 \times \text{time}$ ,  $\text{O}_3 \times \text{time}$  and  $\text{CO}_2 \times \text{community} \times \text{time}$  interactions (Tables 3, 4). This variation in community responses to the treatments is not surprising given the



**Fig. 1** Stand-level biomass of young forest stands exposed to a factorial arrangement of atmospheric  $\text{CO}_2$  and tropospheric  $\text{O}_3$  treatments for 7 yr at the Aspen FACE project in Rhinelander, WI, USA. Values are means ( $n = 3$ ); bars are SEM total biomass. Treatment indicators: 1, control (ambient  $\text{CO}_2$ , ambient  $\text{O}_3$ ); 2, elevated  $\text{CO}_2$  (target  $560 \mu\text{l l}^{-1}$ ); 3, elevated  $\text{O}_3$  ( $1.5\times$  ambient); 4, combined (elevated  $\text{CO}_2 + \text{O}_3$ ).

**Table 3** Beginning (1997) and end (2003) biomass pools (g biomass m<sup>-2</sup>) and yearly biomass production (g biomass m<sup>-2</sup> yr<sup>-1</sup>) by community and atmospheric treatment for foliage, wood (stems + branches), and coarse roots (tap root + all roots >1.0 mm diameter)

Parameter	Control			Elevated CO <sub>2</sub>			Elevated O <sub>3</sub>			+CO <sub>2</sub> , +O <sub>3</sub>		
	AA	AB	AM	AA	AB	AM	AA	AB	AM	AA	AB	AM
Pool size, 1997												
Foliage	1.5 (0.1)	1.4 (0.3)	0.9 (0.0)	1.6 (0.1)	1.5 (0.0)	1.0 (0.1)	1.5 (0.2)	1.4 (0.1)	0.8 (0.0)	1.5 (0.1)	1.4 (0.1)	1.0 (0.0)
Wood	7.6 (0.4)	5.7 (1.0)	2.5 (0.1)	8.0 (0.2)	6.1 (0.1)	3.2 (0.3)	7.3 (1.1)	6.0 (0.3)	2.5 (0.2)	7.2 (0.6)	5.7 (0.7)	2.8 (0.3)
Coarse roots	11.2 (0.3)	7.2 (0.8)	5.9 (0.1)	12.2 (0.2)	7.4 (0.2)	8.4 (0.5)	11.5 (1.2)	6.8 (0.2)	4.7 (0.3)	11.6 (0.7)	7.0 (0.9)	7.1 (0.4)
Fine roots	0.7 (0.0)	0.5 (0.0)	0.7 (0.0)	0.9 (0.0)	0.5 (0.0)	0.8 (0.0)	1.4 (0.1)	0.4 (0.0)	0.6 (0.0)	1.6 (0.1)	0.6 (0.1)	0.9 (0.0)
Foliage, wood and coarse root production, and fine root biomass, 1998												
Foliage	46.5 (5.8)	38.8 (8.4)	15.7 (1.0)	48.8 (5.0)	51.5 (3.9)	24.2 (3.1)	34.8 (2.6)	33.4 (3.2)	14.4 (1.4)	37.7 (4.1)	37.4 (2.3)	17.7 (5.4)
Wood	226.1 (29.3)	174.8 (38.1)	70.2 (4.7)	237.4 (25.2)	233.1 (18.9)	109.4 (15.0)	166.9 (12.7)	148.4 (13.6)	65.0 (7.8)	181.6 (21.0)	167.4 (9.9)	77.0 (24.7)
Coarse roots	105.0 (9.8)	79.8 (13.4)	41.2 (1.8)	112.7 (10.1)	100.9 (7.8)	71.3 (7.5)	87.9 (3.4)	64.0 (3.8)	31.7 (2.6)	95.1 (8.2)	77.2 (3.5)	46.9 (12.6)
Fine roots	7.8 (0.7)	5.7 (0.9)	5.7 (0.2)	9.5 (0.8)	8.1 (0.6)	7.4 (0.7)	11.9 (0.5)	4.7 (0.2)	4.9 (0.3)	15.2 (1.1)	7.4 (0.3)	7.3 (1.8)
Foliage, wood and coarse root production, and fine root biomass, 1999												
Foliage	139.2 (7.3)	132.1 (14.7)	65.0 (4.6)	169.5 (8.6)	187.8 (4.7)	116.7 (5.5)	99.5 (5.7)	108.5 (9.3)	49.6 (11.8)	107.7 (5.6)	130.2 (9.2)	77.3 (18.9)
Wood	471.6 (11.1)	452.2 (38.0)	236.4 (23.4)	614.7 (38.9)	667.4 (27.8)	465.4 (45.1)	327.1 (16.2)	358.6 (36.1)	173.9 (53.5)	356.1 (8.1)	449.3 (35.4)	297.4 (75.9)
Coarse roots	135.8 (3.0)	131.6 (7.4)	81.6 (6.9)	175.9 (10.7)	181.8 (7.7)	159.4 (14.2)	107.5 (2.2)	101.3 (8.0)	51.4 (10.7)	115.9 (1.4)	130.5 (9.0)	100.4 (18.3)
Fine roots	16.9 (0.6)	14.4 (1.3)	15.7 (0.8)	23.0 (1.0)	21.8 (0.6)	22.2 (0.6)	24.8 (0.8)	11.6 (0.6)	11.8 (1.8)	31.8 (1.2)	18.9 (1.0)	20.9 (4.1)
Foliage, wood and coarse root production, and fine root biomass, 2000												
Foliage	234.7 (11.5)	208.9 (21.3)	110.7 (7.8)	301.4 (12.0)	311.3 (21.0)	166.0 (5.9)	168.1 (6.0)	180.0 (9.5)	90.0 (23.7)	196.9 (9.8)	216.5 (14.9)	135.6 (29.3)
Wood	390.5 (24.4)	260.0 (13.4)	216.3 (21.0)	538.9 (34.9)	443.4 (71.7)	250.6 (38.9)	278.8 (12.4)	260.0 (22.3)	194.7 (62.2)	376.5 (45.1)	317.4 (54.1)	292.5 (70.0)
Coarse roots	106.2 (5.8)	90.3 (6.7)	60.6 (4.9)	144.8 (7.8)	134.5 (21.0)	67.9 (12.8)	88.3 (3.1)	79.3 (3.2)	45.7 (8.5)	112.1 (10.2)	100.0 (14.9)	75.2 (16.6)
Fine roots	24.1 (0.9)	20.4 (1.7)	23.1 (1.3)	34.0 (1.3)	31.9 (2.1)	28.5 (0.6)	35.4 (0.5)	16.9 (0.5)	18.0 (2.9)	47.8 (1.7)	27.7 (1.5)	31.1 (5.6)
Foliage, wood and coarse root production, and fine root biomass, 2001												
Foliage	271.3 (11.2)	244.5 (19.8)	152.0 (11.3)	376.9 (22.1)	393.7 (45.1)	244.1 (14.3)	194.2 (7.6)	217.9 (13.7)	125.0 (32.7)	252.6 (19.2)	289.1 (35.5)	188.5 (38.3)
Wood	209.0 (12.1)	191.7 (16.6)	193.9 (21.8)	404.1 (62.9)	450.0 (118.9)	386.0 (48.7)	167.9 (7.4)	207.3 (18.1)	167.9 (47.5)	306.9 (54.6)	372.1 (126.9)	264.4 (61.3)
Coarse roots	35.1 (2.0)	37.8 (2.2)	49.8 (3.2)	70.0 (12.1)	83.2 (24.5)	96.9 (6.3)	25.7 (1.3)	38.2 (4.2)	35.6 (6.3)	57.4 (9.5)	74.2 (24.7)	58.3 (10.6)
Fine roots	26.4 (0.8)	22.9 (1.6)	29.2 (1.6)	39.4 (2.0)	38.2 (3.9)	37.5 (1.2)	38.5 (0.7)	19.5 (0.8)	22.8 (3.7)	56.0 (2.9)	34.2 (3.2)	38.9 (6.4)
Foliage, wood and coarse root production, and fine root biomass, 2002												
Foliage	355.1 (15.8)	295.9 (24.8)	187.6 (10.9)	507.8 (32.9)	496.5 (45.7)	299.2 (16.2)	246.3 (12.5)	265.9 (10.6)	164.8 (44.2)	335.1 (27.7)	346.1 (32.7)	237.6 (49.6)
Wood	434.5 (28.6)	296.4 (7.4)	177.7 (17.0)	695.4 (73.2)	567.0 (55.6)	295.8 (37.3)	283.1 (30.3)	252.2 (29.2)	193.3 (59.5)	431.4 (61.3)	376.0 (47.8)	244.2 (54.4)
Coarse roots	77.8 (5.4)	52.5 (4.3)	38.8 (3.5)	113.4 (7.5)	93.7 (7.3)	64.4 (5.5)	51.0 (3.6)	46.9 (8.4)	36.6 (8.2)	80.9 (10.6)	55.1 (8.0)	50.7 (11.6)
Fine roots	31.7 (1.1)	26.4 (1.8)	34.0 (1.5)	48.1 (2.5)	45.3 (4.1)	43.5 (1.5)	44.6 (1.0)	22.6 (0.5)	27.8 (4.6)	67.6 (3.8)	39.1 (2.9)	45.8 (7.8)
Foliage, wood and coarse root production, and fine root biomass, 2003												
Foliage	450.8 (17.4)	360.2 (26.7)	232.7 (12.9)	627.8 (49.0)	600.5 (48.5)	365.1 (21.3)	320.8 (19.1)	329.8 (12.7)	213.5 (58.7)	428.5 (30.4)	452.9 (43.2)	295.2 (62.7)
Wood	500.1 (9.8)	333.0 (9.8)	244.6 (52.3)	642.7 (89.5)	568.9 (51.7)	352.2 (48.7)	405.9 (28.6)	325.7 (19.5)	242.7 (75.7)	484.9 (17.7)	534.6 (60.0)	295.4 (69.1)
Coarse roots	82.3 (1.6)	62.1 (2.6)	42.1 (3.7)	91.5 (10.5)	86.1 (12.9)	68.6 (5.3)	66.6 (5.3)	60.1 (3.3)	39.7 (8.9)	84.5 (1.7)	101.8 (9.1)	53.9 (13.3)
Fine roots	37.2 (1.2)	30.5 (1.9)	39.1 (1.7)	55.1 (3.3)	51.7 (4.1)	50.0 (1.8)	52.5 (1.6)	26.6 (0.6)	33.2 (5.7)	79.7 (3.8)	48.0 (3.6)	53.1 (9.5)
Pool size, 2003*												
Foliage	450.8 (17.4)	360.2 (26.7)	232.7 (12.9)	627.8 (49.0)	600.5 (48.5)	365.1 (21.3)	320.8 (19.1)	329.8 (12.7)	213.5 (58.7)	428.5 (30.4)	452.9 (43.2)	295.2 (62.7)
Wood	2307.3 (89.4)	1773.9 (132.1)	1125.2 (67.4)	3223.2 (253.4)	3003.4 (246.1)	1816.7 (108.4)	1638.8 (99.2)	1611.9 (63.4)	1026.2 (306.1)	2192.8 (157.3)	2246.4 (216.2)	1461.1 (323.6)
Coarse roots	553.5 (18.4)	461.3 (29.5)	320.1 (13.8)	720.6 (43.9)	687.7 (54.7)	537.1 (19.6)	438.6 (13.6)	396.6 (9.4)	245.6 (42.5)	557.4 (26.4)	545.9 (41.7)	392.6 (70.6)
Fine roots	37.2 (1.2)	30.5 (1.9)	39.1 (1.7)	55.1 (3.3)	51.7 (4.1)	50.0 (1.8)	52.5 (1.6)	26.6 (0.6)	33.2 (5.7)	79.7 (3.8)	48.0 (3.6)	53.1 (9.5)

Fine root (<1.0 mm diameter) data are standing crop at height of growing season (mid-July). Values are means (n = 3) with standard error in parentheses. Community indicators: AA, pure aspen; AB, aspen–birch mix; AM, aspen–maple mix.

\*Wood and coarse root pool size estimates for 2003 were determined using allometric biomass equations and may differ slightly from estimates arrived at by summing annual production caused by rounding. Differences are generally <3% and within the standard error of the allometric estimates.



**Table 4** Statistical significance of atmospheric CO<sub>2</sub> (CO<sub>2</sub>), tropospheric O<sub>3</sub> (O<sub>3</sub>) and community experimental factors on stand-level foliage, wood and coarse root production from 1998 to 2003 at the Aspen FACE project in Rhinelander, WI, USA

Source	Foliage	Wood	Coarse roots
Block	ns	ns	ns
CO <sub>2</sub>	0.041	0.009	0.004
O <sub>3</sub>	0.039	0.039	0.017
CO <sub>2</sub> × O <sub>3</sub>	ns	ns	ns
Community	0.001	ns	ns
CO <sub>2</sub> × community	ns	ns	ns
O <sub>3</sub> × community	ns	0.057	ns
CO <sub>2</sub> × O <sub>3</sub> × community	ns	ns	ns
Time	<0.000	<0.000	<0.000
CO <sub>2</sub> × time	<0.000	<0.000	0.001
O <sub>3</sub> × time	0.001	0.002	<0.000
CO <sub>2</sub> × O <sub>3</sub> × time	ns	ns	ns
Community × time	<0.000	<0.000	<0.000
CO <sub>2</sub> × community × time	ns	0.059	0.031
O <sub>3</sub> × community × time	ns	ns	ns
CO <sub>2</sub> × O <sub>3</sub> × community × time	ns	ns	ns

Data were analyzed using a repeated-measures split-plot ANOVA after King *et al.* (2001).

Wood, stem + branches; coarse roots, roots >1 mm diameter; ns, not statistically significant ( $P > 0.05$ ).

Fine root production was not determined in this study.

interannual variability of weather, host-specific pathogens, etc. in such a long-term experiment. On average, elevated CO<sub>2</sub> increased foliage, wood and coarse root production from 31 to 71% relative to the control, with the aspen–birch and aspen–maple communities showing larger relative responses than pure aspen (Table 5). Elevated O<sub>3</sub> decreased annual production of foliage, wood and coarse roots relative to the control by 9–29% on average, with the aspen community generally showing the largest decrease in growth. Concurrent exposure to elevated CO<sub>2</sub> and O<sub>3</sub> caused small reductions in aspen foliage, wood and coarse root annual production relative to the control, and 15–30% increases in the aspen–birch and aspen–maple communities (Tables 4, 5).

Relative to the control, average July fine root biomass increased 45, 64 and 29% under elevated CO<sub>2</sub> in the aspen, aspen–birch and aspen maple communities, respectively,

Tables 3, 5, 6). Tropospheric ozone caused an average decrease in fine root biomass relative to the control of 15 and 19% in the aspen–birch and aspen–maple communities (Tables 3, 5, 6), respectively, but curiously increased it 44% in the aspen community. In the +CO<sub>2</sub>, +O<sub>3</sub> treatment, mean July fine root biomass increased relative to the control 107, 46 and 34% in the aspen, aspen–birch and aspen–maple communities, respectively (Tables 3, 5, 6).

### Biomass accumulation

Changes in annual production caused by the experimental treatments during the 7-yr experimental period had cumulative impacts on standing biomass in the aspen, aspen–birch and aspen–maple communities (Table 6; Fig. 1). By 2003, total tree biomass in control plots averaged 3349, 2626 and 1717 g m<sup>-2</sup> in the aspen, aspen–birch and aspen–maple communities, respectively (Table 3; Fig. 1). Biomass was distributed on average as 13.8% foliage, 67.3% wood (stems + branches), 17.6% coarse roots, and 1.5% in fine roots, with minor variations between communities. Allometric analyses (King *et al.*, 1996) showed that shifts between the major biomass fractions because of the treatments were statistically insignificant (data not shown).

Relative to the control, elevated CO<sub>2</sub> increased total biomass by 24.9, 45.6 and 60.3% averaged over all years in aspen, aspen–birch and aspen–maple communities, respectively (Fig. 2). The degree of stimulation increased over time, with a significant CO<sub>2</sub> × time interaction (Table 6; Fig. 2). Tropospheric O<sub>3</sub> decreased standing biomass relative to the control by an average 22, 13 and 14% in the aspen, aspen–birch and aspen–maple communities, respectively (Table 6; Fig. 2). The aspen community sustained the largest decline, whereas growth in the aspen–birch and aspen–maple communities became less sensitive to tropospheric O<sub>3</sub> over time (Table 6; Fig. 2). Concurrent exposure to elevated CO<sub>2</sub> provided some protection against the negative effects of O<sub>3</sub> (Figs 1, 2). Standing biomass in the aspen community under the combined treatment (+CO<sub>2</sub>, +O<sub>3</sub>) declined on average 8% relative to the control, but by the fourth year of treatment had recovered to near ambient levels. Biomass production in aspen–birch and aspen–maple communities under concurrent exposure was

Parameter	+CO <sub>2</sub>			+O <sub>3</sub>			+CO <sub>2</sub> + O <sub>3</sub>		
	AA	AB	AM	AA	AB	AM	AA	AB	AM
Foliage	35.7	59.4	59.1	-29.0	-11.3	-13.9	-9.3	15.0	24.6
Wood	40.4	71.5	63.2	-27.0	-9.1	-8.9	-4.2	29.8	29.1
Coarse roots	30.6	49.8	68.3	-21.2	-14.2	-23.4	0.7	18.7	22.7
Fine roots	45.1	63.8	28.8	44.1	-15.3	-19.3	106.9	45.7	34.3

**Table 5** Average relative response (percentage change relative to control) of foliage, wood (stems + branches), and coarse root mean annual production, and mean fine root biomass from 1998 to 2003, to experimental treatments at the Aspen FACE project in Rhinelander, WI, USA

Values calculated as (treatment mean – control mean)/control mean × 100.

Community indicators: AA, pure aspen; AB, aspen–birch mix; AM, aspen–maple mix.

**Table 6** Statistical significance of atmospheric CO<sub>2</sub>, tropospheric O<sub>3</sub> and community experimental factors on stand-level biomass from 1997 to 2003 at the Aspen FACE project in Rhinelander, WI, USA

Source	Foliage	Wood	Coarse roots	Fine roots	Total
Block	ns	ns	ns	ns	ns
CO <sub>2</sub>	0.041	0.008	0.004	0.003	0.008
O <sub>3</sub>	0.039	0.015	0.008	ns	0.016
CO <sub>2</sub> × O <sub>3</sub>	ns	ns	ns	ns	ns
Community	0.001	0.016	ns	<0.000	0.031
CO <sub>2</sub> × community	ns	ns	ns	0.031	ns
O <sub>3</sub> × community	ns	0.009	ns	<0.000	0.043
CO <sub>2</sub> × O <sub>3</sub> × community	ns	ns	ns	ns	ns
Time	<0.000	<0.000	<0.000	<0.000	<0.000
CO <sub>2</sub> × time	<0.000	<0.000	<0.000	<0.000	<0.000
O <sub>3</sub> × time	0.001	<0.000	<0.000	ns	<0.000
CO <sub>2</sub> × O <sub>3</sub> × time	ns	ns	ns	ns	ns
Community × time	<0.000	<0.000	<0.000	<0.000	<0.000
CO <sub>2</sub> × community × time	ns	ns	ns	0.011	ns
O <sub>3</sub> × community × time	ns	<0.000	ns	0.036	0.001
CO <sub>2</sub> × O <sub>3</sub> × community × time	ns	ns	ns	ns	ns

Data were analyzed using a repeated-measures split-plot ANOVA after King *et al.* (2001). Wood, stems + branches; coarse roots, all roots >1 mm diameter; fine roots, roots <1 mm diameter; ns, not statistically significant ( $P > 0.05$ ).

comparable with, or greater than, that in control plots, with a stimulation of 8 and 24%, respectively, averaged over the 7-yr period (Fig. 2).

## Discussion

The Aspen FACE project has provided a unique experimental platform for investigating forest ecosystem responses to the rising concentrations of atmospheric CO<sub>2</sub> and tropospheric O<sub>3</sub>. A hallmark of the experiment is the consistency of response of many ecosystem properties to the atmospheric treatments (Karnosky *et al.*, 2003, 2005), including growth (Isebrands *et al.*, 2001; Percy *et al.*, 2002; Karnosky *et al.*, 2005); leaf physiology (Noormets *et al.*, 2001; Takeuchi *et al.*, 2001); soil respiration and soil C cycling (King *et al.*, 2001, 2004; Loya *et al.*, 2003; Karberg *et al.*, 2005); and soil N transformations and microbial dynamics (Larson *et al.*, 2002; Holmes *et al.*, 2003; Zak *et al.*, 2003). The primary driver of many of these ecosystem-level responses to the experimental treatments is NPP. We hypothesized that NPP would be stimulated by elevated atmospheric CO<sub>2</sub> and decreased by tropospheric O<sub>3</sub>. Combined fumigation was expected to result in NPP similar to that of the control. This analysis of 7 yr of NPP data largely supports these hypotheses.


### Allometric modeling of biomass and NPP

Growth of the experimental forest communities of the Aspen FACE project compares well with growth of stands of similar age reported from sites in north-central Wisconsin, Minnesota and Alaska. After 7 yr growth, control stands of pure aspen at our site had total above-ground (wood + foliage) and below-ground (coarse + fine roots) biomass of

2758 and 591 g m<sup>-2</sup>, respectively. Ruark & Bockheim (1988) reported above- and below-ground biomass in naturally regenerated 8-yr-old aspen stands in north-central Wisconsin of 2500 and 1380 g m<sup>-2</sup>, respectively. The discrepancy in below-ground biomass was probably caused by residual root biomass from the previous stand in the study of Ruark & Bockheim (1988), whereas in our study cuttings were planted in root-free soil.

Alban & Perala (1990) reported average above-ground biomass of 2376 g m<sup>-2</sup> from a series of sites 5 yr after harvest in northern Michigan and Minnesota. Paré & Van Cleve (1993) reported above-ground biomass of naturally regenerated aspen 14 yr after harvest near Fairbanks, Alaska of *c.* 5000 g m<sup>-2</sup>, roughly twice the age and biomass of our study. The agreement between our study and these published reports gives confidence that the Aspen FACE stands are representative of natural forests, and supports our allometric approach to estimating biomass.

Applying species- and site-specific biomass regressions to annual nondestructive measurements of all trees in the experimental plots at Aspen FACE will be valid for estimating biomass and NPP of wood and coarse roots for some time to come (perhaps with some additional destructive harvesting). However, the utility of the allometric approach for fine roots may be limited. There has been difficulty in developing stand-level scaling relationships between above-ground plant parts and soil core estimates of fine root biomass (Kurz *et al.*, 1996; King *et al.*, 1999). This is because of high spatial variation in root distributions, and extremely plastic fine root responses to differences in environmental conditions (Nadelhoffer, 2000; King *et al.*, 2002; Pregitzer, 2002). In the current study, we partitioned a fraction of heart root biomass (determined allometrically) to fine roots based on measured root biomass



**Fig. 2** Response of total biomass production of young forest stands exposed to a factorial arrangement of atmospheric CO<sub>2</sub> and tropospheric O<sub>3</sub> treatments for 7 yr at the Aspen FACE project in Rhinelander, WI, USA. Calculated as (treatment biomass – control biomass)/control biomass × 100.

partitioning from the destructive harvests in 2000 and 2002, of which the fine root sampling was much more rigorous in 2002. The fraction of total root biomass partitioned to fine roots varies with seasonal changes in fine root standing crop (Hendrick & Pregitzer, 1996; King *et al.*, 2002) and stage of stand development (King *et al.*, 1999). Hence, fine root biomass determined from our 'static' partitioning may or may not accurately reflect standing fine root biomass at a given point in time, and does not capture fine root production and turnover.

Similarly, allometric modeling of foliage biomass (NPP) at Aspen FACE has been valid for the early stage of stand development, but its utility will be limited in the future. Comparison of allometric estimates of stand-level foliage production (this study) with litter-trap estimates of litter production showed good agreement ( $R^2 = 0.89$ ) for the years 2001–3

(C. P. Giardina, unpublished). However, foliage biomass becomes 'uncoupled' from stem growth after canopy closure in young stands, and is inversely related to it in older forests (Ovington, 1957; Ford, 1984; Cannell, 1985; Gower *et al.*, 1994; Miller, 1995). Collection of annual foliar litter production suggests the aspen and aspen-birch stands are approaching canopy closure; that is, the annual increment in litter production is decreasing (C.P.G., unpublished data). Therefore future stem growth may not be accompanied by proportional increments in foliage biomass, compromising the allometric approach.

#### Stimulation of NPP by elevated CO<sub>2</sub>

This analysis of 7 yr of growth data supports our first hypothesis, that elevated atmospheric CO<sub>2</sub> (*c.* 550 ppm by



volume) will cause sustained enhancement of forest NPP and biomass accumulation. This is consistent with the thousands of studies conducted at smaller spatial and temporal scales and across a wide variety of plant species over the past several decades (Strain & Bazzaz, 1983; Eamus & Jarvis, 1989; Ceulemans & Mousseau, 1994; Strain & Cure, 1994; Amthor, 1995; Curtis & Wang, 1998; Norby *et al.*, 1999). Stimulation of total biomass accumulation by elevated atmospheric CO<sub>2</sub> at Aspen FACE averaged 43% for all communities, although this response developed over time (significant CO<sub>2</sub> × time interaction). This is higher than the average 25% growth enhancement reported for other forest FACE experiments (DeLucia *et al.*, 1999; Hamilton *et al.*, 2002; Norby *et al.*, 2002; Calfapietra *et al.*, 2003), and the average 31% stimulation from a meta-analysis of the earlier elevated CO<sub>2</sub> literature (Curtis & Wang, 1998).

The large, sustained CO<sub>2</sub> enhancement of NPP at Aspen FACE could have several causes. Because of the relatively high latitude of the site (45°40' N), the soil is of recent glacial origin, with good chemical and physical properties for forest growth (Dickson *et al.*, 2000). Analysis of soil N cycling from 1999 to 2003 (Holmes *et al.*, 2003, 2005) suggests that soil N availability is not constraining growth responses to elevated CO<sub>2</sub> at the Aspen FACE experiment. The north-temperate climate is mesic, with favorable site water balance for most of the year, because of low evaporative demand (calculated by King *et al.*, 2001). A large fraction of the global forest C sink occurs in recently glaciated north-temperate and boreal forest ecosystems, where long-term C storage in soils is especially important (Schlesinger, 1997; Myneni *et al.*, 2001). Therefore the nutrient limitation to sustained CO<sub>2</sub> enhancement of forest NPP, as reported from low-latitude forests on highly weathered soils (Oren *et al.*, 2001), may be less of a constraint at higher latitudes.

Additionally, the experimental stands at Aspen FACE are dominated by early successional species in the early stage of stand development, which confers greater growing space (less intertree competition) and greater productivity relative to older stands (Pregitzer & Euskirchen, 2004). This could provide the capacity for greater stimulation of NPP and other ecosystem properties in response to elevated CO<sub>2</sub> relative to older, closed-canopy forests (King *et al.*, 2004).

The stimulation of total biomass production at Aspen FACE was caused by proportional increases in all plant parts: roots, wood and foliage. Averaged across community type from 1997 to 2003, elevated CO<sub>2</sub> caused 42, 45 and 41% increases in foliage, wood and coarse root biomass, respectively. Allometric analyses on an individual tree basis, using the harvest data from 2000 and 2002 and at the stand level, showed that elevated CO<sub>2</sub> did not change biomass partitioning among plant parts (data not shown). This is consistent with our understanding of tree biomass partitioning responses to elevated atmospheric CO<sub>2</sub> (Gebauer *et al.*, 1996; King *et al.*, 1996; Norby *et al.*, 1999).

There were important differences in the magnitude of CO<sub>2</sub> enhancement of component and total plant biomass production between communities, however. The order of relative response was generally pure aspen < aspen–paper birch < aspen–sugar maple; however, the order of absolute stand-level biomass production has been pure aspen > aspen–paper birch > aspen–sugar maple. In the pure aspen community, it is possible that intraspecific competition has constrained the potential stand-level relative growth enhancement in response to elevated atmospheric CO<sub>2</sub>. Interspecific competition in the aspen–birch community could possibly have allowed a greater overall growth response to elevated CO<sub>2</sub>. The aspen–maple community started out with smaller trees and therefore less intertree competition (intra- and interspecific competition was reduced), hence there was a greater capacity to respond to elevated atmospheric CO<sub>2</sub>.

McDonald *et al.* (2002) provide evidence that competitively advantaged trees in the pure aspen community at Aspen FACE show a greater relative growth response to elevated CO<sub>2</sub> compared with competitively disadvantaged trees in an autoregressive manner ('the big get bigger faster'). These results apparently scale to the level of the stand. It will be interesting to see how relative growth responses to the treatments change as the stands proceed through canopy closure, and intertree competition and mortality become more significant. More growing space in the young stands could contribute to the greater relative CO<sub>2</sub> growth enhancement at Aspen FACE compared with the Duke (DeLucia *et al.*, 1999; Hamilton *et al.*, 2002) and Oak Ridge National Laboratory (Norby *et al.*, 2002) experiments, which both have older, closed canopy forests. This is consistent with a recent synthesis of soil respiration results from the four forest FACE experiments, which found that the relative stimulation of soil respiration caused by elevated CO<sub>2</sub> was greater in young, open-canopy forests compared with older, closed-canopy forests (King *et al.*, 2004).

### Decreased forest NPP from tropospheric O<sub>3</sub>

Our second hypothesis was that elevated tropospheric O<sub>3</sub> (*c.* 1.5 ×) would decrease forest NPP, which was again supported by this analysis of 7 yr of growth data from the Aspen FACE experiment. This result is largely consistent with the literature but, importantly, we feel provides realistic quantification of the magnitude of the response for an important forest type in north-temperate and boreal forest ecosystems. Our understanding of O<sub>3</sub> effects on vegetation is largely based on studies of crops or small trees grown in highly controlled environments for short periods (reviewed by Heck *et al.*, 1984; Pye, 1988; Samuelson & Kelly, 2001; Andersen, 2003). These studies show that, in a wide range of plant species, tropospheric O<sub>3</sub> causes almost universal reductions in crop yield or biomass production, but the magnitude of response has been highly dependent on experimental conditions. High

variation in experimental results and uncertain correlation between visible foliar injury and yield reduction have led to considerable efforts to compare seedling responses with those of mature trees to determine appropriate factors for scaling  $O_3$  responses to the landscape (Chappelka & Samuelson, 1998; Matyssek & Innes, 1999; Samuelson & Kelly, 2001).

At Aspen FACE, chronic exposure to moderately elevated tropospheric  $O_3$  (*c.* 1.5 $\times$ ) has resulted in an average reduction in biomass production of 22, 12 and 16% in the pure aspen, aspen–birch and aspen–maple communities, respectively. These results are comparable with the average 23% decrease in tree seedling growth reported in the review of Pye (1988), but higher than the 2.6–6.8% decrease in annual NPP in the USA during the late 1980s to early 1990s from the modeling study of Felzer *et al.* (2004). Importantly, the response to  $O_3$  was modified by both community composition and time (significant  $O_3 \times$  community  $\times$  time interaction). The pure aspen community was the most sensitive to  $O_3$  and maintained this sensitivity over time. The aspen–birch and aspen–maple communities, however, appear to be losing sensitivity to  $O_3$  relative to the control.

Differences in community response could be caused by compensatory growth of less- $O_3$ -sensitive species (Pye, 1988; Broadmeadow & Jackson, 2000) in the mixed communities, or changes in  $O_3$  responsiveness induced by competition (McDonald *et al.*, 2002; Liu *et al.*, 2004). In the aspen–maple community, sugar maple comprises *c.* 9% of wood biomass, whereas in the aspen–birch community the two species are more evenly represented with no clear dominance of one over the other (data not shown). Hence compensatory growth of less- $O_3$ -sensitive species is unlikely to be the cause of the increased productivity over time. In a 2-yr phytotron study, Liu *et al.* (2004) observed that European beech experienced no reduction in total biomass production caused by elevated  $O_3$  when grown in monoculture. However, when grown in mixed culture with Norway spruce  $O_3$  caused a 32% reduction in beech biomass, and the spruce benefited (+13%) from the weak performance of its competitor. These results underscore the important fact that monospecific responses to  $O_3$  are not simply additive, and more realistic experimental designs are required to determine long-term ecosystem responses to the changing atmosphere. An important aspect of the Aspen FACE experiment will be to see if the mixed communities fully regain productive capacity under elevated  $O_3$ .

As with elevated  $CO_2$ , growth under tropospheric  $O_3$  does not appear to have altered biomass partitioning among the major plant parts, as there were no statistically significant shifts in root : shoot, foliage : branch or wood : coarse root ratios (data not shown). This finding apparently contradicts many earlier studies that show relative decreases in root growth under  $O_3$ , which is thought to aid in the repair of damaged photosynthetic structures by increased C allocation above ground (Karnosky *et al.*, 1996; Andersen, 2003). In our

study, all plant parts became proportionally smaller under elevated  $O_3$ . The exception to this is fine roots in the pure aspen section, which showed an average 44% stimulation in biomass. Possible causes include (i) spurious values among the three replicates for each treatment; (ii) confounding by abundant fine roots from herbaceous species that proliferated under the open canopies of the elevated  $O_3$  treatment; or (iii) it is a real effect. Fine root (<1-mm-diameter) biomass values for each replicate of the 2002 harvest upon which the static partitioning was based were 50.4, 48.1 and 34.0 g m<sup>-2</sup>, compared with an average 31.7 g m<sup>-2</sup> for control plots at that time. If herbaceous roots were accidentally included in our estimates, the static partitioning used here would propagate the error through each year of biomass estimation. This is unlikely as all communities were harvested and processed at the same time using the same method. Sampling error could also have biased towards high root biomass estimates, but this is unlikely as 10 cores of 15 cm diameter  $\times$  25 cm deep were used in each split-plot FACE ring section. The only other fine root harvest at the site performed in 1999 did not detect significant effects of  $O_3$  on fine root biomass (King *et al.*, 2001). In any case, this finding is highly counterintuitive, and requires further study before we can conclude that elevated  $O_3$  increases fine root biomass in young aspen ecosystems.

#### Tropospheric $O_3$ compromises stimulation of NPP caused by elevated $CO_2$

Our final hypothesis was that chronic exposure to elevated  $CO_2$  and elevated tropospheric  $O_3$  ( $+CO_2, +O_3$ ) would result in forest NPP similar to that of the control. A putative 'protective effect' of elevated  $CO_2$  has been discussed (Allen, 1990; Wüstman *et al.*, 2003), in that decreased stomatal conductance under elevated  $CO_2$  might decrease the flux of  $O_3$  into the plant; there may be other protective mechanisms, such as responses of antioxidant enzymes (Rao *et al.*, 1995). Our analysis partially supports this hypothesis. For total and component biomass production, the interaction between  $CO_2$  and  $O_3$  was never statistically significant. Thus elevated  $CO_2$  provided comparable stimulation to NPP at both levels of the  $O_3$  treatment in all communities over time. Because of the sensitivity of the aspen community to  $O_3$ , however, total biomass production in this community was depressed for the first 3 yr of growth, after which it did not differ from the control. The aspen–birch and aspen–maple communities exhibited an average stimulation of total biomass production of 8 and 24%, respectively, under combined fumigation. Thus the large stimulation in biomass production all three communities experienced in response to elevated  $CO_2$  was completely annulled or greatly reduced by concurrent exposure to moderate levels of tropospheric  $O_3$ .

Experiments using long-term exposure of trees to combined  $CO_2$  and  $O_3$  fumigation are beginning to show that responses to both gases are variable, depending on species/clone and

provenance. However, the antagonistic effects on growth of elevated CO<sub>2</sub> and tropospheric O<sub>3</sub> have generally been observed in these experiments. Dickson *et al.* (1998) exposed five hybrid poplar genotypes to factorial treatments of CO<sub>2</sub> and O<sub>3</sub> in open-top chambers (OTC) for 1 yr, and found that plants exposed to combined fumigation (+CO<sub>2</sub>, +O<sub>3</sub>) had biomass similar to the control. Broadmeadow & Jackson (2000) grew seedlings of oak, ash and pine under factorial CO<sub>2</sub> and O<sub>3</sub> treatments in OTC for 3 yr. Elevated CO<sub>2</sub> enhanced growth; O<sub>3</sub> decreased it; and combined fumigation provided some protection from O<sub>3</sub> in the order of species responsiveness: oak > pine > ash. In a 5-yr OTC experiment, Rebbeck & Scherzer (2002) found that yellow poplar growth was insensitive to O<sub>3</sub> alone, but increased 60% with combined fumigation (+CO<sub>2</sub>, +O<sub>3</sub>) relative to the control, but not until the fifth season. Similarly, Riikonen *et al.* (2004) found that growth of two clones of silver birch responded negatively to O<sub>3</sub>, but only at ambient CO<sub>2</sub>. Tree growth increased under elevated CO<sub>2</sub> and combined fumigation (+CO<sub>2</sub>, +O<sub>3</sub>) treatments. Together with our results these studies show that, in the long run, elevated CO<sub>2</sub> provides some protection from exposure to phytotoxic concentrations of tropospheric O<sub>3</sub> for a variety of forest tree species. However, this also means that gains in NPP that could be achieved under elevated CO<sub>2</sub> are being compromised by tropospheric O<sub>3</sub> pollution, and this has had continent-scale implications for C sequestration for some time (Felzer *et al.*, 2004).

Because CO<sub>2</sub> is chemically inert in the atmosphere, and human population growth and fossil energy consumption continue to increase, the concentration of atmospheric CO<sub>2</sub> will continue to rise for the foreseeable future. Tropospheric O<sub>3</sub> is highly reactive and was historically considered a regional pollutant. However, it is becoming apparent that the cumulative impact of industrialization around the globe is also raising the background concentration of this pollutant along with CO<sub>2</sub>. Our analysis of 7 yr of growth data at the Aspen FACE project, and decades of earlier research, indicate that the concentration of atmospheric CO<sub>2</sub> expected for the year 2050 has the capacity to stimulate forest NPP. At least for young northern forests on glacial soils, this response does not appear to be constrained by nutrient or water limitations. However, the concurrent global rise in tropospheric O<sub>3</sub> is damaging forest physiology and growth to the point that potential gains in terrestrial C sequestration caused by rising CO<sub>2</sub> are partially or completely annulled. We conclude, therefore, that global monitoring of ambient O<sub>3</sub> exposure of vegetation should become an important part of government environmental protection programs. Moving forward with technologies that remove important anthropogenic precursors to photochemical O<sub>3</sub> formation (mainly oxidized forms of nitrogen) from automobile and industrial emissions would help to decrease concentrations of tropospheric O<sub>3</sub> because of its short half-life in the atmosphere, decreasing at least one constraint on the capacity of forest ecosystems to sequester atmospheric C.

## Acknowledgements

Aspen FACE is principally supported by the Office of Science (BER), US Department of Energy, Grant No. DE-FG02-95ER62125 to Michigan Technological University, and Contract No. DE-AC02-98CH10886 to Brookhaven National Laboratory, the US Forest Service Northern Global Change Program and North-central Research Station, Michigan Technological University, and Natural Resources Canada – Canadian Forest Service. Additional support for research at Aspen FACE was supported by DOE Program of Ecosystem Research (PER) grant number DE-FG02-93ER6166, the USDA Forest Service Northern Global Change Program, and USDA NRI Competitive Grants Program nos. 2001-35107-11262 and 2004-35102-14782. The authors thank Jaak Sober and Wendy Jones for their assistance in operating the Aspen FACE facility.

## References

- Adams RM, Hamilton SA, McCarl BA. 1986. The benefits of pollution control: the case for ozone and US agriculture. *American Journal of Agricultural Economics* 68: 886–893.
- Alban DH, Perala DA. 1990. Ecosystem carbon following aspen harvesting in the Upper Great Lakes. In: Adams RD, ed. *Aspen Symposium' 89. Proceedings*. General Technical Report NC-140. St Paul, MN, USA: USDA Forest Service, 123–131.
- Allen JH. 1990. Plant response to rising carbon dioxide and potential interactions with air pollutants. *Journal of Environmental Quality* 19: 15–34.
- Amthor JS. 1995. Terrestrial higher-plant response to increasing atmospheric [CO<sub>2</sub>] in relation to the global carbon cycle. *Global Change Biology* 1: 243–274.
- Andersen CP. 2003. Source–sink balance and carbon allocation below ground in plants exposed to ozone. *New Phytologist* 157: 213–228.
- Baskerville GL. 1972. Use of logarithmic regression in the estimation of plant biomass. *Canadian Journal of Forest Research* 2: 49–53.
- Birdsey RA, Lewis GM. 2002. Current and historical trends in use, management, and disturbance of US forestlands. In: Kimble JM, Heath LS, Birdsey RA, Lal R, eds. *The Potential of US Forest Soils to Sequester Carbon and Mitigate the Greenhouse Effect*. Boca Raton, FL, USA: CRC Press, 15–33.
- Broadmeadow MSJ, Jackson SB. 2000. Growth responses of *Quercus petraea*, *Fraxinus excelsior* and *Pinus sylvestris* to elevated carbon dioxide, ozone and water supply. *New Phytologist* 146: 437–451.
- Calfapietra C, Gielen B, Galema ANJ, Lukac M, DeAngelis P, Moscatelli MC, Ceulemans R, Scarascia-Mugnozza G. 2003. Free-air CO<sub>2</sub> enrichment (FACE) enhances biomass production in a short-rotation poplar plantation. *Tree Physiology* 23: 805–814.
- Cannell MGR. 1985. Dry matter partitioning in tree crops. In: Cannell MGR, Jackson JE, eds. *Attributes of Trees as Crop Plants*. Huntingdon, UK: Natural Environmental Research Council, 160–193.
- Caspersen JP, Pacala SW, Jenkins JC, Hurtt GC, Moorcroft PR, Birdsey RA. 2000. Contributions of land-use history to carbon accumulation in US forests. *Science* 290: 1148–1151.
- Ceulemans R, Mousseau M. 1994. Effects of elevated atmospheric CO<sub>2</sub> on woody plants. *New Phytologist* 127: 425–446.
- Chappelka AH, Samuelson LJ. 1998. Ambient ozone effects on forest trees of the eastern United States: a review. *New Phytologist* 139: 91–108.
- Curtis PS, Wang X. 1998. A meta-analysis of elevated CO<sub>2</sub> effects on woody plant mass, form, and physiology. *Oecologia* 113: 299–313.



- DeLucia EH, Hamilton JG, Naidu SL, Thomas RB, Andrews JA, Finzi A, Lavine M, Matamala R, Mohan JE, Hendrey GR, Schlesinger WH. 1999. Net primary production of a forest ecosystem with experimental CO<sub>2</sub> enrichment. *Science* 284: 1177–1179.
- Dickson RE, Coleman MD, Riemenschneider DE, Isebrands JG, Hogan GD, Karnosky DF. 1998. Growth of five hybrid poplar genotypes exposed to interacting elevated CO<sub>2</sub> and O<sub>3</sub>. *Canadian Journal of Forest Research* 28: 1706–1716.
- Dickson RE, Lewin KF, Isebrands JG, Coleman MD, Heilman WE, Riemenschneider DE, Sober J, Host GE, Zak DR, Hendrey GR, Pregitzer KS, Karnosky DF. 2000. *Forest Atmosphere Carbon Transfer and Storage (FACTS-II): The Aspen Free-air CO<sub>2</sub> and O<sub>3</sub> Enrichment (FACE) Project: An Overview*. General Technical Report NC-214. St Paul, MN, USA: USDA Forest Service NCRS.
- Eamus D, Jarvis PG. 1989. The direct effects of increase in the global atmospheric CO<sub>2</sub> concentration on natural and commercial temperate forests. *Advances in Ecological Research* 19: 1–55.
- Felzer B, Kicklighter D, Mellilo J, Wang C, Zhuang Q, Prinn R. 2004. Effects of ozone on net primary production and carbon sequestration in the conterminous United States using a biogeochemistry model. *Tellus* 56B: 230–248.
- Ford ED. 1984. The dynamics of plantation growth. In: Bowen GD, Nambiar EKS, eds. *Nutrition of Plantation Forests*. London, UK: Academic Press, 17–52.
- Fowler D, Cape JN, Coyle M, Flechard C, Kuylenstierna J, Hicks K, Derwent D, Johnson C, Stevenson D. 1999. The global exposure of forests to air pollutants. *Water, Air and Soil Pollution* 116: 5–32.
- Gebauer RLE, Reynolds JF, Strain BR. 1996. Allometric relations and growth in *Pinus taeda*: the effect of elevated CO<sub>2</sub> and changing N availability. *New Phytologist* 134: 85–93.
- Gower ST, Gholz HL, Nakane K, Baldwin VC. 1994. Production and carbon allocation patterns of pine forests. *Ecological Bulletins* 43: 115–135.
- Hamilton JG, DeLucia EH, George K, Naidu SL, Finzi AC, Schlesinger WH. 2002. Forest carbon balance under elevated CO<sub>2</sub>. *Oecologia* 131: 250–260.
- Heck WW, Cure WW, Rawlings JO, Zaragoza LJ, Heagle AS, Heggstad HE, Kohut RJ, Kress LW, Temple PJ. 1984. Assessing impacts of ozone on agricultural crops. I. Overview. *Journal of the Air Pollution Control Association* 34: 729–735.
- Hendrey GR, Ellsworth DS, Lewin KF, Nagy J. 1999. A free-air enrichment system for exposing tall forest vegetation to elevated atmospheric CO<sub>2</sub>. *Global Change Biology* 5: 293–309.
- Hendrick RL, Pregitzer KS. 1996. Temporal and depth-related patterns of fine root dynamics in northern hardwood forests. *Journal of Ecology* 84: 167–176.
- Holmes WE, Zak DR, Pregitzer KS, King JS. 2003. Soil nitrogen transformations under *Populus tremuloides*, *Betula papyrifera* and *Acer saccharum* following 3 years exposure to elevated CO<sub>2</sub> and O<sub>3</sub>. *Global Change Biology* 9: 1743–1750.
- Holmes WE, Zak DR, Pregitzer KS, King JS. 2005. Elevated CO<sub>2</sub> and O<sub>3</sub> alter soil nitrogen transformations beneath trembling aspen, paper birch, and sugar maple. *Ecosystems*. (in press.)
- Houghton JT, Ding Y, Griggs DJ, Noguer M, van der Linden PJ, Dai X, Maskell K, Johnson CA. 2001. *Climate Change 2001: The Scientific Basis*. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change. New York, NY, USA: Cambridge University Press.
- Houghton RA, Hackler JL, Lawrence KT. 1999. The US carbon budget: contributions from land-use change. *Science* 285: 574–578.
- Isebrands JG, McDonald EP, Kruger E, Hendrey G, Pregitzer K, Percy K, Söber J, Karnosky DF. 2001. Growth responses of *Populus tremuloides* clones to interacting carbon dioxide and tropospheric ozone. *Environmental Pollution* 115: 359–371.
- Johnson DW. 1999. Simulated nitrogen cycling response to elevated CO<sub>2</sub> in *Pinus taeda* and mixed deciduous forests. *Tree Physiology* 19: 321–327.
- Karberg NJ, Pregitzer KS, King JS, Friend AL, Wood JR. 2005. Soil carbon dioxide partial pressure and dissolved inorganic carbonate chemistry under elevated carbon dioxide and ozone. *Oecologia* 142: 296–306.
- Karnosky DF, Gagnon ZE, Dickson RE, Coleman MD, Lee EH, Isebrands JG. 1996. Changes in growth, leaf abscission, and biomass associated with seasonal tropospheric ozone exposures of *Populus tremuloides* clones and seedlings. *Canadian Journal of Forest Research* 26: 23–37.
- Karnosky DF, Mankovska B, Percy K, Dickson RE, Podila DG, Sober J, Noormets A, Hendrey GR, Coleman MD, Kubiske M, Pregitzer KS, Isebrands JG. 1999. Effects of tropospheric O<sub>3</sub> on trembling aspen and interaction with CO<sub>2</sub>: results from an O<sub>3</sub>-gradient and a FACE experiment. *Water, Air, and Soil Pollution* 116: 311–322.
- Karnosky DF, Zak DR, Pregitzer KS, Awmack CS, Bockheim JG, Dickson RE, Hendrey GR, Host GE, King JS, Kopper BJ, Kruger EL, Kubiske ME, Lindroth RL, Mattson WJ, McDonald EP, Noormets A, Oksanen E, Parsons WFJ, Percy KE, Podila GK, Riemenschneider DE, Sharma P, Thakur R, Söber J, Jones WS, Anttonen S, Vapaavuori E, Mankovska B, Heilman W, Isebrands JG. 2003. Tropospheric O<sub>3</sub> moderates responses of temperate hardwood forests to elevated CO<sub>2</sub>: a synthesis of molecular to ecosystem results from the Aspen FACE project. *Functional Ecology* 17: 289–304.
- Karnosky DF, Pregitzer KS, Zak DR, Kubiske ME, Hendrey GR, Weinstein D, Nosal M, Percy KE. 2005. Scaling ozone responses of forest trees to the ecosystem level in a changing climate. *Plant, Cell & Environment* 28: 965–981.
- King JS, Thomas RB, Strain BR. 1996. Growth and carbon accumulation in root systems of *Pinus taeda* and *Pinus ponderosa* seedlings as affected by varying CO<sub>2</sub>, temperature, and nitrogen. *Tree Physiology* 16: 635–642.
- King JS, Albaugh TJ, Allen LH, Kress LW. 1999. Stand-level allometry in *Pinus taeda* as affected by irrigation and fertilization. *Tree Physiology* 19: 769–778.
- King JS, Pregitzer KS, Zak DR, Sober J, Isebrands JG, Dickson RE, Hendrey GR, Karnosky DF. 2001. Fine-root biomass and fluxes of soil carbon in young stands of paper birch and trembling aspen as affected by elevated atmospheric CO<sub>2</sub> and tropospheric O<sub>3</sub>. *Oecologia* 128: 237–250.
- King JS, Albaugh TJ, Allen LH, Buford M, Strain BR, Dougherty P. 2002. Below-ground carbon input to soil is controlled by nutrient availability and fine root dynamics in loblolly pine. *New Phytologist* 154: 389–398.
- King JS, Hanson PJ, Bernhardt E, DeAngelis P, Norby RJ, Pregitzer KS. 2004. A multi-year synthesis to elevated atmospheric CO<sub>2</sub> from four forest FACE experiments. *Global Change Biology* 10: 1027–1042.
- Kopp RJ, Vaughn WJ, Hazilla M, Carson R. 1985. Implications of environmental policy for US agriculture: the case of ambient ozone standards. *Journal of Environmental Management* 20: 321–331.
- Krupa S, McGrath MT, Andersen CP, Booker FL, Burkey KO, Chappelka AH, Chevone BI, Pell EJ, Zilinskas BA. 2000. Ozone and plant health. *Plant Disease* 85: 4–12.
- Kurz WA, Beukema SJ, Apps MJ. 1996. Estimation of root biomass and dynamics for the carbon budget model of the Canadian forest sector. *Canadian Journal of Forest Research* 26: 1973–1979.
- Larson J, Zak DR, Sinsabaugh RL. 2002. Extracellular enzyme activity and metabolism of root-derived substrates beneath temperate trees growing under elevated CO<sub>2</sub> and O<sub>3</sub>. *Soil Science Society of America Journal* 66: 1848–1856.
- Levy HII, Kasibhatla PS, Moxim WJ, Klonecki AA, Hirsch AI, Oltmans SJ, Chameides WL. 1997. The global impact of human activity on tropospheric ozone. *Geophysical Research Letters* 24: 791–794.
- Liu X, Kozovits AR, Grams TEE, Blaschke H, Rennenberg H, Matyssek R. 2004. Competition modifies effects of enhanced ozone/carbon dioxide concentrations on carbohydrate and biomass accumulation in juvenile Norway spruce and European beech. *Tree Physiology* 24: 1045–1055.
- Loya WM, Pregitzer KS, Karberg NJ, King JS, Giardina CP. 2003. Reduction of soil carbon formation by tropospheric ozone under increased carbon dioxide levels. *Nature* 425: 705–707.

- Matussek R, Innes JL. 1999. Ozone – a risk factor for trees and forests in Europe? *Water, Air, and Soil Pollution* 116: 199–226.
- McDonald EP, Kruger EL, Riemenschneider DE, Isebrands JG. 2002. Competitive status influences tree-growth responses to elevated CO<sub>2</sub> and O<sub>3</sub> in aggrading stands. *Functional Ecology* 16: 792–801.
- McGuire AD, Melillo JM, Joyce LA. 1995. The role of nitrogen in the response of forest net primary production to elevated atmospheric carbon dioxide. *Annual Review of Ecology and Systematics* 26: 473–503.
- Miller HG. 1995. The influence of stand development on nutrient demand, growth and allocation. *Plant Soil* 168–169: 225–232.
- Murphy JJ, Deluki MA, McCubbin DR, Kim HJ. 1999. The cost of crop damage caused by ozone air pollution from motor vehicles. *Journal of Environmental Management* 55: 273–289.
- Myneni RB, Dong J, Tucker CJ, Kaufman RK, Kauppi PE, Liski J, Zhou L, Alexeyev V, Hughes MK. 2001. A large carbon sink in the woody biomass of Northern forests. *Proceedings of the National Academy of Sciences, USA* 98: 14784–14789.
- Nadelhoffer KJ. 2000. The potential effects of nitrogen deposition on fine root production in forest ecosystems. *New Phytologist* 147: 131–139.
- Noormets A, Sober A, Pell EJ, Dickson RE, Podila GK, Sober J, Isebrands JG, Karnosky DF. 2001. Stomatal and non-stomatal limitation to photosynthesis in two trembling aspen (*Populus tremuloides* Michx.) clones exposed to elevated CO<sub>2</sub> and/or O<sub>3</sub>. *Plant, Cell & Environment* 24: 327–336.
- Norby RJ, Wullschlegel SD, Gunderson CA, Johnson DW, Ceulemans R. 1999. Tree responses to rising CO<sub>2</sub> in field experiments: implications for the future forest. *Plant, Cell & Environment* 22: 683–714.
- Norby RJ, Hanson PJ, O'Neill EG, Tschaplinski TJ, Weltzin JF, Hansen RA, Cheng W, Wullschlegel SD, Gunderson CA, Edwards NT, Johnson DW. 2002. Net primary production of a CO<sub>2</sub>-enriched deciduous forest and the implications for carbon storage. *Ecological Applications* 12: 1261–1266.
- Oren R, Ellsworth DS, Johnsen KH, Phillips N, Ewers BE, Maier C, Schafer KVR, McCarthy H, Hendrey G, McNulty SG, Katul GG. 2001. Soil fertility limits carbon sequestration by forest ecosystems in a CO<sub>2</sub>-enriched atmosphere. *Nature* 411: 469–472.
- Ovington JD. 1957. Dry-matter production by *Pinus sylvestris* L. *Annals of Botany* 21: 287–314.
- Pacala SW, Hurtt GC, Baker D, Peylin P, Houghton RA, Birdsey RA, Heath L, Sundquist ET, Stallard RF, Ciais P, Moorcroft P, Caspersen JP, Shevliakova E, Moore B, Kohlmaier G, Holland E, Gloor M, Harmon ME, Fan S-M, Sarmiento JL, Goodale CL, Schimel D, Field CB. 2001. Consistent land- and atmosphere-based US carbon sink estimates. *Science* 292: 2316–2319.
- Paré D, Van Cleve K. 1993. Aboveground biomass production and nutrient accumulation on postharvested white spruce sites in interior Alaska. *Canadian Journal of Forest Research* 23: 1233–1239.
- Percy KE, Awmack CS, Lindroth RL, Kubiske ME, Kopper BJ, Isebrands JG, Pregitzer KS, Hendrey GR, Dickson RE, Zak DR, Oksanen E, Sober J, Harrington R, Karnosky DF. 2002. Altered performance of forest pests under atmospheres enriched by CO<sub>2</sub> and O<sub>3</sub>. *Nature* 420: 403–407.
- Pregitzer KS. 2002. Fine roots of trees – a new perspective. *New Phytologist* 154: 267–273.
- Pregitzer KS, Euskirchen ES. 2004. Carbon cycling and storage in world forests: biome patterns related to forest age. *Global Change Biology* 10: 2052–2077.
- Pye JM. 1988. Impact of ozone on the growth and yield of trees: a review. *Journal of Environmental Quality* 17: 347–360.
- Rao MV, Hale BA, Ormrod DP. 1995. Amelioration of ozone-induced oxidative damage in wheat plants grown under high carbon dioxide. *Plant Physiology* 109: 421–432.
- Rebbeck J, Scherzer AJ. 2002. Growth responses of yellow poplar (*Liriodendron tulipifera* L.) exposed to 5 years of O<sub>3</sub> alone or combined with elevated CO<sub>2</sub>. *Plant, Cell & Environment* 25: 1527–1537.
- Riikonen J, Lindsberg M-M, Holopainen T, Oksanen E, Lappi J, Peltonen P, Vapaavuori E. 2004. Silver birch and climate change: variable growth and carbon allocation responses to elevated concentrations of carbon dioxide and ozone. *Tree Physiology* 24: 1227–1237.
- Ruark GA, Bockheim JG. 1988. Biomass, net primary production, and nutrient distribution for an age sequence of *Populus tremuloides* ecosystems. *Canadian Journal of Forest Research* 18: 435–443.
- Samuelson L, Kelly JM. 2001. Scaling ozone effects from seedlings to forest trees. *New Phytologist* 149: 21–41.
- Schlesinger WH. 1997. *Biogeochemistry: An Analysis of Global Change*. San Diego, CA, USA: Academic Press.
- Strain BR, Bazzaz FA. 1983. Terrestrial plant communities. In: Lemon ER, ed. *CO<sub>2</sub> and Plants: The Response of Plants to Rising Levels of Atmospheric Carbon Dioxide*. Boulder, CO, USA: Westview Press, 177–222.
- Strain BR, Cure JD. 1994. *Direct Effects of Atmospheric CO<sub>2</sub> Enrichment on Plants and Ecosystems: An Updated Bibliographic Database*. ORNL/CDIAC-70. Oak Ridge, TN, USA: Oak Ridge National Laboratory.
- Takeuchi Y, Kubiske ME, Isebrands JG, Pregitzer KS, Hendrey G, Karnosky DF. 2001. Photosynthesis, light and nitrogen relationships in a young deciduous forest canopy under open-air CO<sub>2</sub> enrichment. *Plant, Cell & Environment* 24: 1257–1268.
- Wustman BA, Oksanen E, Karnosky DF, Noormets A, Isebrands JG, Pregitzer KS, Hendrey GR, Sober J, Podila GK. 2003. Effects of elevated CO<sub>2</sub> and O<sub>3</sub> on aspen clones of varying O<sub>3</sub> sensitivity. In: Karnosky DF, Percy KE, Chappelka AH, Simpson C, Pikkariainen J, eds. *Air Pollution, Global Change and Forests in the New Millennium*. Boston, MA, USA: Elsevier, 391–410.
- Zak DR, Holmes WE, Finzi AC, Norby RJ, Schlesinger WH. 2003. Soil nitrogen cycling under elevated CO<sub>2</sub>: a synthesis of forest FACE experiments. *Ecological Applications* 13: 1508–1514.
- Zak DR, Pregitzer KS, Curtis PS, Vogel CS, Holmes WE, Lussenhop J. 2000. Atmospheric CO<sub>2</sub>, soil-N availability, and allocation of biomass and nitrogen by *Populus tremuloides*. *Ecological Applications* 10: 34–46.





### About *New Phytologist*

- *New Phytologist* is owned by a non-profit-making **charitable trust** dedicated to the promotion of plant science, facilitating projects from symposia to open access for our Tansley reviews. Complete information is available at [www.newphytologist.org](http://www.newphytologist.org).
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as-ready' via *OnlineEarly* – the 2004 average submission to decision time was just 30 days. Online-only colour is **free**, and essential print colour costs will be met if necessary. We also provide 25 offprints as well as a PDF for each article.
- For online summaries and ToC alerts, go to the website and click on 'Journal online'. You can take out a **personal subscription** to the journal for a fraction of the institutional price. Rates start at £109 in Europe/\$202 in the USA & Canada for the online edition (click on 'Subscribe' at the website).
- If you have any questions, do get in touch with Central Office ([newphytol@lancaster.ac.uk](mailto:newphytol@lancaster.ac.uk); tel +44 1524 594691) or, for a local contact in North America, the US Office ([newphytol@ornl.gov](mailto:newphytol@ornl.gov); tel +1 865 576 5261).